



The second eigenvalue of the fractional p -Laplacian

Lorenzo Brasco, Enea Parini

► To cite this version:

Lorenzo Brasco, Enea Parini. The second eigenvalue of the fractional p -Laplacian. *Advances in Calculus of Variation*, 2016, 10.1515/acv-2015-0007 . hal-01066698

HAL Id: hal-01066698

<https://hal.science/hal-01066698>

Submitted on 22 Sep 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright

THE SECOND EIGENVALUE OF THE FRACTIONAL p -LAPLACIAN

LORENZO BRASCO AND ENEA PARINI

ABSTRACT. We consider the eigenvalue problem for the *fractional p -Laplacian* in an open bounded, possibly disconnected set $\Omega \subset \mathbb{R}^n$, under homogeneous Dirichlet boundary conditions. After discussing some regularity issues for eigenfunctions, we show that the second eigenvalue $\lambda_2(\Omega)$ is well-defined, and we characterize it by means of several equivalent variational formulations. In particular, we extend the mountain pass characterization of Cuesta, De Figueiredo and Gossez to the nonlocal and nonlinear setting. Finally, we consider the minimization problem

$$\inf\{\lambda_2(\Omega) : |\Omega| = c\}.$$

We prove that, differently from the local case, an optimal shape does not exist, even among disconnected sets. A minimizing sequence is given by the union of two disjoint balls of volume $c/2$ whose mutual distance tends to infinity.

CONTENTS

1. Introduction	2
1.1. Overview and aim of the paper	2
1.2. Results of the paper	3
1.3. Some words about regularity	4
1.4. Plan of the paper	5
2. Definitions and preliminary results	5
2.1. Notation	5
2.2. Sobolev spaces	5
2.3. Nonlocal eigenvalues	7
3. A journey into regularity for the fractional p -Laplacian	9
3.1. Global boundedness	9
3.2. Caccioppoli inequalities	12
3.3. Local L^∞ estimates	17
3.4. Basic comparison estimates	21
3.5. A theorem by Kuusi, Mingione and Sire	24
4. The second eigenvalue	25
5. Mountain pass characterization	29
6. A sharp lower bound	32
Appendix A. Some useful inequalities I	34
Appendix B. Some useful inequalities II	36
References	38

Date: September 19, 2014.

2010 Mathematics Subject Classification. 35P30, 47J10, 35R09.

Key words and phrases. Nonlocal eigenvalue problems, spectral optimization, quasilinear nonlocal operators, Caccioppoli estimates.

1. INTRODUCTION

1.1. Overview and aim of the paper. Let $\Omega \subset \mathbb{R}^N$ be a bounded open set, $0 < s < 1$ and $1 < p < \infty$. This paper is concerned with the nonlinear and nonlocal Dirichlet eigenvalue problem

$$(1.1) \quad (-\Delta_p)^s u = \lambda |u|^{p-2} u, \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \mathbb{R}^N \setminus \Omega,$$

where

$$(-\Delta_p)^s u(x) := 2 \lim_{\delta \rightarrow 0^+} \int_{\{y \in \mathbb{R}^N : |y-x| \geq \delta\}} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} dy,$$

is the *fractional p -Laplacian*. Here solutions of (1.1) are always understood in the weak sense, see equation (2.7) below.

If $\lambda \in \mathbb{R}$ is such that (1.1) admits a solution $u \not\equiv 0$, then we say that λ is an (s, p) -*eigenvalue* of Ω . Correspondingly, u is an (s, p) -*eigenfunction* associated to λ . The eigenvalue problem (1.1) was first introduced by Lindgren and Lindqvist in [21] and investigated by several authors afterwards, we cite for example [4, 14] and [18].

Observe that for $p = 2$ the operator $(-\Delta_p)^s$ reduces to the well-known fractional Laplacian, which has been extensively studied in the last years (see for example [5, 15] and the references therein). We point out that the terminology *fractional p -Laplacian* is not standard, though somehow justified by the fact that the operator $(-\Delta_p)^s$ arises as the first variation of the fractional Dirichlet integral

$$(1.2) \quad \Phi_{s,p}(u) = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy,$$

and therefore it is the nonlocal counterpart of the usual p -Laplacian operator

$$-\Delta_p u := -\operatorname{div}(|\nabla u|^{p-2} \nabla u).$$

By homogeneity, it is not difficult to see that (s, p) -eigenvalues corresponds to critical points of the functional (1.2) restricted to the manifold $\mathcal{S}_p(\Omega)$ consisting of functions having unitary L^p -norm.

In order to put the studies of this paper in the right framework, let us start by recalling some known facts about the eigenvalue problem (1.1). First of all, in analogy with the local case, the *spectrum of the fractional p -Laplacian*, i.e the set

$$\sigma_{s,p}(\Omega) = \{\lambda \in \mathbb{R} : \lambda \text{ is an } (s, p)\text{-eigenvalue}\}$$

is a closed set (see [21]). Moreover, it is possible to define the first eigenvalue $\lambda_1(\Omega)$, i.e. the smallest $\lambda \in \sigma_{s,p}(\Omega)$. The first eigenvalue has a variational characterization, as it corresponds to the minimum of $\Phi_{s,p}$ on $\mathcal{S}_p(\Omega)$. In other words, $\lambda_1(\Omega)$ coincides with the sharp constant in the following Poincaré inequality (see [4, Lemma 2.4])

$$c \int_{\Omega} |u|^p dx \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy, \quad u \in C_0^\infty(\Omega).$$

The quantity $\lambda_1(\Omega)$ can be estimated from below in terms of $|\Omega|$ in a sharp way, exactly as in the local case. This is a consequence of the *Faber-Krahn inequality* (see [4, Theorem 3.5])

$$\lambda_1(\Omega) \geq \left(\frac{|B|}{|\Omega|} \right)^{\frac{sp}{N}} \lambda_1(B),$$

where B is any N -dimensional ball. Equality in the previous holds if and only if Ω itself is a ball. The result can also be rephrased by saying:

“Among all domains of fixed volume, the ball has the smallest first eigenvalue.”

The main aim of this paper is to study the *second eigenvalue* $\lambda_2(\Omega)$ of the fractional p -Laplacian, for every $1 < p < \infty$ and $0 < s < 1$. Observe that since we are dealing with a *nonlinear* eigenvalue problem, actually it is not even clear that it is possible to speak about a second eigenvalue. Indeed, the spectrum $\sigma_{s,p}(\Omega)$ in principle could contain a sequence accumulating to $\lambda_1(\Omega)$.

Thus, at first we want to show that $\lambda_2(\Omega)$ is well-defined and give a variational characterization for it. Then we want to prove sharp lower bounds for $\lambda_2(\Omega)$ in terms of $|\Omega|$, similarly to the Faber-Krahn inequality. It is useful to recall at this point that in the local case this is indeed possible, thanks to the *Hong-Krahn-Szego inequality*. This asserts that:

“Among all domains of fixed volume, the disjoint union of two equal balls has the smallest second eigenvalue.”

In scaling invariant form, this reads as

$$\lambda_2(\Omega) \geq \left(\frac{2|B|}{|\Omega|} \right)^{\frac{p}{N}} \lambda_1(B),$$

where B is again any N -dimensional ball. Equality holds if and only if Ω itself is a disjoint union of two equal balls. For $p = 2$ the result was proved long time ago by Krahn [19]. Then this has been probably neglected and rediscovered some years later by Hong [17] and P. Szego [23]. The general case of the p -Laplacian has been recently addressed in [3, Theorem 3.2].

1.2. Results of the paper. The first main result of this paper is the following (see Sections 4 and 5 for the precise statements).

The second eigenvalue. *Let $1 < p < \infty$ and $0 < s < 1$. Let $\Omega \subset \mathbb{R}^N$ be an open and bounded set. There exists a real positive number $\lambda_2(\Omega)$ with the following properties:*

- $\lambda_2(\Omega)$ is an (s, p) -eigenvalue;
- $\lambda_2(\Omega) > \lambda_1(\Omega)$;
- if $\lambda > \lambda_1(\Omega)$ is an (s, p) -eigenvalue, then $\lambda \geq \lambda_2(\Omega)$.

Moreover, it has the following variational characterization

$$(1.3) \quad \lambda_2(\Omega) = \inf_{\gamma \in \Gamma(u_1, -u_1)} \max_{u \in \text{Im}(\gamma)} \Phi_{s,p}(u),$$

where $\Gamma(u_1, -u_1)$ is the set of continuous paths on $\mathcal{S}_p(\Omega)$ connecting the first eigenfunction u_1 to its opposite $-u_1$.

The eigenvalue $\lambda_2(\Omega)$ will be constructed by means of a variational minimax procedure originally introduced by Drábek and Robinson in [12]. In particular, in this paper we will deliberately avoid to use any index theory. The mountain pass characterization (1.3) is the nonlocal counterpart of the result by Cuesta, De Figueiredo and Gossez for the local case, see [7]. We point out that our proof differs from that of [7] and is similar to that of [2, Proposition 5.4], which is based on the so called *hidden convexity* for Dirichlet integrals (see [1]).

On the contrary for the Hong-Krahn-Szego inequality the situation in the local case and in the nonlocal are quite different. Due to the nonlocal effects, the mutual position of the connected components of the domain influences the spectrum $\sigma_{s,p}(\Omega)$ of the operator. More important, as already observed in [21], if u is a sign-changing eigenfunction with eigenvalue λ and Ω_+ is one of its nodal domains, *it is no more true that $\lambda_1(\Omega_+) = \lambda$* (see Lemma 6.1 below). Also, the restriction of u to Ω_+ in general is not a first eigenfunction of Ω_+ . *This point marks a huge difference with the local case.* We then have the following sharp estimate for $\lambda_2(\Omega)$, which is partially in contrast with the local case. This is the second main result of the paper.

Nonlocal Hong-Krahn-Szego inequality (Nonlocal Hong-Krahn-Szego inequality). *Let $1 < p < \infty$ and $0 < s < 1$. For every $\Omega \subset \mathbb{R}^N$ open and bounded set we have*

$$(1.4) \quad \lambda_2(\Omega) > \left(\frac{2|B|}{|\Omega|} \right)^{\frac{s p}{N}} \lambda_1(B),$$

where B is any N -dimensional ball. Equality is never attained in (1.4), but the estimate is sharp in the following sense: if Ω_n is a disjoint union of two equal balls $B_R(x_n)$ and $B_R(y_n)$ such that

$$\lim_{n \rightarrow \infty} |x_n - y_n| = +\infty,$$

then

$$\lim_{n \rightarrow \infty} \lambda_2(\Omega_n) = \left(\frac{|B|}{|B_R|} \right)^{\frac{s p}{N}} \lambda_1(B) = \lambda_1(B_R).$$

Observe that as a consequence of this result, we obtain that the shape optimization problem

$$\inf \{ \lambda_2(\Omega) : |\Omega| = c \},$$

does not admit a solution.

1.3. Some words about regularity. Our proofs are based on variational techniques for which membership of eigenfunctions to the relevant Sobolev space is sufficient.

The only place where regularity of eigenfunctions is really needed in our paper is in the proof of the nonlocal Hong-Krahn-Szego inequality, which is based on the crucial Lemma 6.1. There continuity is necessary in order to assure that nodal domains of eigenfunctions are open sets, thus we enclose in this paper the (long) Section 3 where the regularity issue is tackled. There we prove at first some local and global L^∞ estimates for solutions of general nonhomogeneous equations like

$$(1.5) \quad (-\Delta_p)^s u = F, \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \mathbb{R}^N \setminus \Omega.$$

Though not completely new¹, these estimates are here obtained by means of a *Moser's iteration technique*. For this reason, we believe them to be interesting and we prefer to include them here.

Then we show how continuity follows from the very recent result [20, Theorem 1.5] by Kuusi, Mingione and Sire. This point needs a precision: the regularity estimates of [20] are indeed very general. In particular, they consider more general nonlinear and nonlocal operators and cover the harder case of F being just a measure. In this case solutions have to be understood in a suitable very weak sense (see [20, Definition 2]). However, such a general setting needs the hypothesis

$$p > 2 - s/N,$$

thus their result can not be directly applied to our situation where $1 < p < \infty$ and $0 < s < 1$ without any further restriction. Such a restriction on p and s in [20] comes from a couple of crucial comparison results (essentially [20, Lemma 3.4]). These results can be proved in an easier way when F is in the correct Lebesgue space, without any additional condition on p and s . The scope of the second part of Section 3 is exactly that of showing how to fix this technical point. Then we briefly sketch the main ideas of the proof of [20, Theorem 1.5] for the reader's convenience.

¹In [8, Theorem 1.1] the local L^∞ estimate is proved for solutions of the homogeneous equation, while [14, Theorem 3.2] contains the global L^∞ bound for eigenfunctions. Both these results use a suitable variant of De Giorgi's technique.

1.4. Plan of the paper. All the definitions, notations and preliminary results needed for the sequel are contained in Section 2. In Section 3 we prove some regularity estimates for eigenfunctions and more generally for solutions of (1.5). The second eigenvalue of the fractional p -Laplacian is then introduced and studied in Section 4, while Section 5 contains its mountain pass characterization. Finally, the nonlocal Hong-Krahn-Szego inequality is proved in Section 6. The paper ends with a couple of Appendices containing some pointwise inequalities needed throughout the whole paper.

Acknowledgements. Part of this work has been conducted during the conference “*Journées d’Analyse Appliquée Nice-Toulon-Marseille*” held in Porquerolles in May 2014. Organizers and hosting institutions are gratefully acknowledged.

2. DEFINITIONS AND PRELIMINARY RESULTS

2.1. Notation. Throughout the whole paper, we will denote by $B_R(x)$ the N -dimensional ball having radius R and center x . When the center will be clear from the context or unnecessary, we will simply write B_R . Finally, ω_N is the measure of the N -dimensional ball with unit radius. For a Borel set $E \subset \mathbb{R}^N$, we will denote by $|E|$ its N -dimensional Lebesgue measure. The average over $B_R(x_0)$ of a measurable function ψ will be denoted by

$$(2.1) \quad \bar{\psi}_{x_0, R} := \frac{1}{|B_R(x_0)|} \int_{B_R(x_0)} \psi \, dx.$$

We also set

$$\psi_+(x) = \max\{\psi(x), 0\} \quad \text{and} \quad \psi_-(x) = \max\{-\psi(x), 0\},$$

so that $\psi = \psi_+ - \psi_-$. Given $1 < p < \infty$ and $0 < s < 1$ such that $sp < N$, we define

$$p^* = \frac{Np}{N-sp} \quad \text{and} \quad (p^*)' = \frac{Np}{Np - N + sp}.$$

2.2. Sobolev spaces. Let $0 < s < 1$ and $1 < p < \infty$. For every $\Omega \subset \mathbb{R}^N$ open and bounded set, we consider the Sobolev space $\widetilde{W}_0^{s,p}(\Omega)$ defined as the closure of $C_0^\infty(\Omega)$ with respect to the norm

$$\|u\|_{\widetilde{W}_0^{s,p}(\Omega)} := \left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} \, dx \, dy \right)^{\frac{1}{p}}, \quad u \in C_0^\infty(\Omega).$$

We recall that the space $\widetilde{W}_0^{s,p}(\Omega)$ can be equivalently defined by taking the closure of $C_0^\infty(\Omega)$ with respect to the full norm

$$\left(\int_{\Omega} |u|^p \, dx \right)^{\frac{1}{p}} + \left(\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} \, dx \, dy \right)^{\frac{1}{p}},$$

see [4, Remark 2.5]. If $sp \neq 1$ and $\partial\Omega$ is smooth enough, such a space coincides with the usual one $W_0^{s,p}(\Omega)$, defined as the closure of $C_0^\infty(\Omega)$ with respect to the norm

$$\left(\int_{\Omega} |u|^p \, dx \right)^{\frac{1}{p}} + \left(\int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} \, dx \, dy \right)^{\frac{1}{p}}.$$

This is proven for example in [4, Proposition B.1].

Remark 2.1 (Borderline case $sp = 1$). When $sp = 1$, the previous identification is no longer true and one has the strict inclusion

$$\widetilde{W}_0^{s,p}(\Omega) \subsetneq W_0^{s,p}(\Omega).$$

Indeed, for $sp \leq 1$ the characteristic function 1_Ω of Ω can be approximated in norm by a sequence $\{u_n\}_{n \in \mathbb{N}} \subset C_0^\infty(\Omega)$, that is we have

$$\lim_{n \rightarrow \infty} \int_\Omega |u_n - 1_\Omega|^p dx = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \int_\Omega \int_\Omega \frac{|u_n(x) - u_n(y)|^p}{|x - y|^{N+sp}} dx dy = 0,$$

see for example the counterexample contained in [11] at page 557. Thus we have $1_\Omega \in W_0^{s,p}(\Omega)$. On the other hand, for $sp = 1$ it is easily seen that

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|1_\Omega(x) - 1_\Omega(y)|^p}{|x - y|^{N+1}} dx dy = 2 \int_\Omega \int_{\mathbb{R}^N \setminus \Omega} \frac{1}{|x - y|^{N+1}} dx dy,$$

and the last integral does not converge, thus $1_\Omega \notin \widetilde{W}_0^{s,p}(\Omega)$.

The main properties of the space $\widetilde{W}_0^{s,p}(\Omega)$ can be found in [4, Section 2]. Here we state and prove a couple of simple functional inequalities that will be needed in the sequel. We use the notation

$$W^{s,p}(\Omega) := \left\{ u \in L^p(\Omega) : \int_\Omega \int_\Omega \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy < +\infty \right\}.$$

Proposition 2.2 (Poincaré with localized seminorm). *Let $1 < p < \infty$ and $0 < s < 1$. We fix $R > 0$, for every $u \in W^{s,p}(B_R)$ we have*

$$(2.2) \quad \frac{|\{x \in B_R : u = 0\}|}{2^{N+sp} R^N} R^{-sp} \int_{B_R} |u|^p dx \leq \int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy.$$

In particular, for $u \in \widetilde{W}_0^{s,p}(B_r)$ with $0 < r < R$ there holds

$$(2.3) \quad \frac{N \omega_N}{2^{N+sp}} \left(\frac{r}{R}\right)^N \left(\frac{R-r}{r}\right) R^{-sp} \int_{B_r} |u|^p dx \leq \int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy.$$

Proof. Let $x \in B_r$ and we pick $y \in B_r$ such that $u(y) = 0$. Then we get

$$|u(x)|^p = |u(x) - u(y)|^p = \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} |x - y|^{N+sp} \leq (2r)^{N+sp} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}}.$$

By integrating at first with respect to y and then integrating with respect to x , we can get inequality (2.2).

In order to prove (2.3), it is sufficient to use (2.2) and observe that

$$|\{x \in B_R : u = 0\}| \geq \omega_N (R^N - r^N).$$

By further noticing that

$$\int_{B_R} |u|^p dx = \int_{B_r} |u|^p dx \quad \text{and} \quad R^N - r^N \geq N r^{N-1} (R - r),$$

we get the conclusion after some simple manipulations. \square

Proposition 2.3 (Sobolev with localized seminorm). *Let $1 < p < \infty$ and $0 < s < 1$ such that $sp < N$. We fix $0 < r < R$, then for every $u \in \widetilde{W}_0^{s,p}(B_r)$ there holds*

$$(2.4) \quad C \left(\int_{B_r} |u|^{p^*} dx \right)^{\frac{p}{p^*}} \leq \int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy,$$

where the constant $C = C(N, p, s, R/r) > 0$ goes to 0 as R/r converges to 1.

Proof. Since $\widetilde{W}_0^{s,p}(B_r) \hookrightarrow W_0^{s,p}(\mathbb{R}^N)$, we have

$$(2.5) \quad \mathcal{S}_{N,p,s} \left(\int_{B_r} |u|^{p^*} dx \right)^{\frac{p}{p^*}} \leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy,$$

see for example [10, Theorem 6.5]. We now decompose the Gagliardo seminorm as

$$\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy = \int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy + 2 \int_{B_R} \int_{\mathbb{R}^N \setminus B_R} \frac{|u(x)|^p}{|x - y|^{N+sp}} dx dy.$$

By observing that $u \equiv 0$ outside B_r and that for every $x \in B_r$ we have $B_{R-r}(x) \subset B_R$, the last integral is easily estimated as follows

$$\begin{aligned} \int_{B_R} \int_{\mathbb{R}^N \setminus B_R} \frac{|u(x)|^p}{|x - y|^{N+sp}} dx dy &= \int_{B_r} \int_{\mathbb{R}^N \setminus B_R} \frac{|u(x)|^p}{|x - y|^{N+sp}} dx dy \\ &\leq \int_{B_r} |u|^p dx \left(\int_{\mathbb{R}^N \setminus B_{R-r}(x_0)} \frac{1}{|y - x_0|^{N+sp}} dy \right) \\ &= \frac{N \omega_N}{s p (R - r)^{sp}} \int_{B_r} |u|^p dx \\ &\leq c_{N,p,s} \left(\frac{R}{r} \right)^N \left(\frac{r}{R - r} \right) \left(\frac{R}{R - r} \right)^{sp} \int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy. \end{aligned}$$

In the last passage we used (2.3). By using the previous in (2.5), we get the conclusion. \square

Remark 2.4 (Watch out!). We point out that for $sp \leq 1$ the previous inequalities (2.3) and (2.4) fail to be true for $R = r$. The counterexample is the same as in Remark 2.1, i.e. the characteristic function of B_R .

Finally, we give a Sobolev inequality without loss on the left-hand side.

Proposition 2.5 (Sobolev with localized full norm). *Let $1 < p < \infty$ and $0 < s < 1$ such that $sp < N$. For every $u \in W^{s,p}(B_R)$ there holds*

$$(2.6) \quad \|u\|_{L^{p^*}(B_R)}^p \leq c \left[\int_{B_R} \int_{B_R} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy + \frac{1}{R^{sp}} \int_{B_R} |u|^p dx \right],$$

for some constant $c = c(N, s, p) > 0$.

Proof. For $R = 1$ the result is contained for example in [10, Theorem 6.7]. The case of a general $R > 0$ can then be obtained by a standard scaling argument. \square

2.3. Nonlocal eigenvalues. We say that $u \in \widetilde{W}_0^{s,p}(\Omega) \setminus \{0\}$ is an (s, p) -eigenfunction associated to the eigenvalue λ if u satisfies (1.1) weakly, i.e.

$$(2.7) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy = \lambda \int_{\Omega} |u|^{p-2} u \varphi dx,$$

for every $\varphi \in \widetilde{W}_0^{s,p}(\Omega)$. If we set

$$\mathcal{S}_p(\Omega) = \left\{ u \in \widetilde{W}_0^{s,p}(\Omega) : \int_{\Omega} |u|^p dx = 1 \right\},$$

we already observed in the Introduction that (s, p) -eigenvalues coincide with critical points of the functional

$$(2.8) \quad \Phi_{s,p}(u) = \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy,$$

restricted to the manifold $\mathcal{S}_p(\Omega)$. The first (s, p) -eigenvalue of Ω is given by

$$(2.9) \quad \lambda_1(\Omega) = \min_{u \in \mathcal{S}_p(\Omega)} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p.$$

It is easy to see that first eigenfunctions must be nonnegative (or nonpositive). This is a consequence of the elementary inequality

$$||u(x)| - |u(y)|| \leq |u(x) - u(y)|,$$

which holds strictly whenever $u(x)u(y) < 0$.

It can be proved actually that any constant sign (s, p) -eigenfunction must be strictly positive (or strictly negative) on every open bounded set Ω , *even disconnected*. This is the content of the following result, which holds true without any connectedness assumptions and which therefore extends [1, Theorem A.1].

Proposition 2.6 (Minimum principle). *Let $1 < p < \infty$ and $0 < s < 1$. Let $u \in \widetilde{W}_0^{s,p}(\Omega) \setminus \{0\}$ be a nonnegative (s, p) -eigenfunction associated to the eigenvalue λ . Then $u > 0$ in Ω .*

Proof. If Ω is connected, this is exactly [1, Theorem A.1]. Let us suppose that Ω is not connected. Still by [1, Theorem A.1] we already know that $u > 0$ on each connected component of Ω where it is not identically zero. Set $\Omega_1 := \{x \in \Omega : u(x) > 0\}$, and suppose by contradiction that there exists a connected component Ω_2 of Ω such that u identically vanishes on Ω_2 . Let $\varphi \in C_0^\infty(\Omega_2)$ be a nonnegative test function, not identically zero. By inserting this in (2.7), with simple manipulations we get

$$\begin{aligned} 0 &= \lambda \int_{\Omega} u^{p-1} \varphi \, dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) \, dx \, dy \\ &= -2 \int_{\Omega_1} \int_{\Omega_2} \frac{u(x)^{p-1}}{|x - y|^{N+sp}} \varphi(y) \, dx \, dy. \end{aligned}$$

Therefore $u \equiv 0$ in Ω_1 , a contradiction. Hence $u > 0$ in Ω . \square

Remark 2.7. The previous result is in contrast with the local case, where actually nonnegative eigenfunctions can identically vanish on some connected components.

The following statement summarizes some basic facts about the first eigenvalue.

Theorem 2.8. *Let $1 < p < \infty$ and $0 < s < 1$. For every $\Omega \subset \mathbb{R}^N$ open and bounded set we have:*

- i) any first (s, p) -eigenfunction must be strictly positive (or strictly negative);*
- ii) $\lambda_1(\Omega)$ is simple, i.e. the solution of (2.9) is unique, up to the choice of the sign;*
- iii) if u is an eigenfunction associated to an eigenvalue $\lambda > \lambda_1(\Omega)$, then u must be sign-changing.*

Proof. The first item follows from Proposition 2.6.

Simplicity of $\lambda_1(\Omega)$ follows from [14, Theorem 4.2], which holds true also for disconnected domains, thanks to Proposition 2.6.

The last statement for general sets follows combining Proposition 2.6 and [14, Theorem 4.1] (again, this last result is valid also without assuming that Ω is connected). \square

Remark 2.9. As in the local case, the exact structure of the spectrum $\sigma_{s,p}(\Omega)$ is an open issue. However, it is possible to show that it certainly contains an increasing sequence of eigenvalues $\{\lambda_k(\Omega)\}_{k \in \mathbb{N}}$ diverging at infinity, by means of standard minimax procedures (see for example [18, Proposition 2.2]). We also mention the interesting result [18, Theorem 1.1] about the asymptotic distribution of (s, p) -eigenvalues.

3. A JOURNEY INTO REGULARITY FOR THE FRACTIONAL p -LAPLACIAN

The aim of this section is to prove that an (s, p) -eigenfunction is continuous, for every $1 < p < \infty$ and $0 < s < 1$. Indeed, we will consider the general case of solutions to

$$(3.1) \quad (-\Delta_p)^s u = F, \quad \text{in } \Omega, \quad u = 0 \quad \text{in } \mathbb{R}^N \setminus \Omega,$$

where $F \in L^{(p^*)}'(\Omega)$ is given. We will prove some global and local L^∞ estimates and then explain how continuity follows from the recent result by Kuusi, Mingione and Sire contained in [20], concerning the case of F being just a measure (not necessarily belonging to the dual of $\widetilde{W}_0^{s,p}(\Omega)$).

We point out that for $sp > N$ we already know that $\widetilde{W}_0^{s,p}(\Omega) \hookrightarrow L^\infty(\Omega) \cap C^{0,s-N/p}(\Omega)$ (see [4, Proposition 2.9]). Then we only need to consider the case $sp \leq N$. For simplicity, we will consider the case $sp < N$, then it will be evident how to handle the borderline case $sp = N$, where $F \in L^q$ for $q > 1$. In this last case, it will be sufficient to reproduce the proofs of this section, by replacing the continuous embedding $W^{s,p} \hookrightarrow L^{p^*}$ (and the associated Sobolev inequality) with $W^{s,N/s} \hookrightarrow L^m$ and $m > 1$ large enough.

Throughout the whole section, given $F \in L^{(p^*)}'(\Omega)$ we always denote by $u \in \widetilde{W}_0^{s,p}(\Omega)$ the solution to (3.1), i.e. u satisfies

$$(3.2) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy = \int_{\Omega} F \varphi dx,$$

for every $\varphi \in \widetilde{W}_0^{s,p}(\Omega)$.

3.1. Global boundedness. We start with the following global result.

Theorem 3.1 (Global L^∞ bound). *Let $1 < p < \infty$ and $0 < s < 1$ such that $sp < N$. If $F \in L^q(\Omega)$ for $q > N/(sp)$, then $u \in L^\infty(\Omega)$. Moreover, we have the scaling invariant estimate*

$$(3.3) \quad \|u\|_{L^\infty(\Omega)} \leq C \left[\frac{1}{|\Omega|} \int_{\Omega} |u| dx + \left(|\Omega|^{\frac{sp}{N} - \frac{1}{q}} \|F\|_{L^q(\Omega)} \right)^{\frac{1}{p-1}} \right],$$

where $C = C(N, p, s, q) > 0$.

Proof. For every $0 < \varepsilon \ll 1$, we define the smooth convex Lipschitz function

$$f_\varepsilon(t) = (\varepsilon^2 + t^2)^{\frac{1}{2}},$$

then we insert the test function $\varphi = \psi |f'_\varepsilon(u)|^{p-2} f'_\varepsilon(u)$ in (3.2), where $\psi \in C_0^\infty(\Omega)$ is a positive function. By using (A.1) with the choices

$$a = u(x), \quad b = u(y), \quad A = \psi(x) \quad \text{and} \quad B = \psi(y),$$

we get

$$\begin{aligned} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|f_\varepsilon(u(x)) - f_\varepsilon(u(y))|^{p-2} (f_\varepsilon(u(x)) - f_\varepsilon(u(y)))}{|x - y|^{N+sp}} (\psi(x) - \psi(y)) dx dy \\ \leq \int_{\Omega} |F| |f'_\varepsilon(u)|^{p-1} \psi dx, \end{aligned}$$

By observing that f_ε converges to $f(t) = |t|$, using that $|f'_\varepsilon(t)| \leq 1$ and Fatou Lemma², if we pass to the limit in the previous we get that the function $|u|$ verifies

$$(3.4) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left| |u(x)| - |u(y)| \right|^{p-2} \left(|u(x)| - |u(y)| \right)}{|x - y|^{N+sp}} (\psi(x) - \psi(y)) dx dy \leq \int_{\Omega} |F| \psi dx,$$

for every positive functions $\psi \in C_0^\infty(\Omega)$. By density the same remains true for $\psi \in \widetilde{W}_0^{s,p}(\Omega)$, still with $\psi \geq 0$.

For every $M > 0$, we define $u_M = \min\{|u|, M\}$ and observe that u_M is still in $\widetilde{W}_0^{s,p}(\Omega)$, since this is just the composition of $|u|$ with a Lipschitz function. Given $\beta > 0$ and $\delta > 0$, we insert the test function $\psi = (u_M + \delta)^\beta - \delta^\beta$ in (3.4), then we get

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left| |u(x)| - |u(y)| \right|^{p-2} \left(|u(x)| - |u(y)| \right) \left((u_M(x) + \delta)^\beta - (u_M(y) + \delta)^\beta \right)}{|x - y|^{N+sp}} dx dy \\ & \leq \int_{\Omega} |F| (u_M + \delta)^\beta dx. \end{aligned}$$

By using inequality (A.3) with the function

$$g(t) = (\min\{t, M\} + \delta)^\beta,$$

from the previous we get

$$\frac{\beta p^p}{(\beta + p - 1)^p} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left| (u_M(x) + \delta)^{\frac{\beta+p-1}{p}} - (u_M(y) + \delta)^{\frac{\beta+p-1}{p}} \right|^p}{|x - y|^{N+sp}} dx dy \leq \int_{\Omega} |F| (u_M + \delta)^\beta dx.$$

We can now use the Sobolev inequality for $W_0^{s,p}(\mathbb{R}^N)$ (see [10, Theorem 6.5]), so to get

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left| (u_M(x) + \delta)^{\frac{\beta+p-1}{p}} - (u_M(y) + \delta)^{\frac{\beta+p-1}{p}} \right|^p}{|x - y|^{N+sp}} dx dy \\ & \geq C_{N,p,s} \left(\int_{\Omega} \left((u_M + \delta)^{\frac{\beta+p-1}{p}} - \delta^{\frac{\beta+p-1}{p}} \right)^{\frac{Np}{N-sp}} dx \right)^{\frac{N-sp}{N}}. \end{aligned}$$

We then obtain

$$\left(\int_{\Omega} \left((u_M + \delta)^{\frac{\beta+p-1}{p}} - \delta^{\frac{\beta+p-1}{p}} \right)^{\frac{Np}{N-sp}} dx \right)^{\frac{N-sp}{N}} \leq \frac{\|F\|_{L^q(\Omega)}}{C_{N,p,s} \beta} \left(\frac{\beta + p - 1}{p} \right)^p \left(\int_{\Omega} (u_M + \delta)^{\beta q'} dx \right)^{\frac{1}{q'}}.$$

We also observe that by triangle inequality and the simple inequality $(u_M + \delta)^{\beta+p-1} \geq \delta^{p-1} (u_M + \delta)^\beta$, the left-hand side can be estimated by

$$\begin{aligned} & \left(\int_{\Omega} \left((u_M + \delta)^{\frac{\beta+p-1}{p}} - \delta^{\frac{\beta+p-1}{p}} \right)^{\frac{Np}{N-sp}} dx \right)^{\frac{N-sp}{N}} \geq \left(\frac{\delta}{2} \right)^{p-1} \left(\int_{\Omega} (u_M + \delta)^\beta \frac{N}{N-sp} dx \right)^{\frac{N-sp}{N}} \\ & \quad - \delta^{\beta+p-1} |\Omega|^{\frac{N-sp}{N}}. \end{aligned}$$

²Observe that the integrand on the left-hand side can be estimated from below by the integrable function

$$-\frac{p-1}{p} \frac{\left| |u(x)| - |u(y)| \right|^p}{|x - y|^{N+sp}} - \frac{1}{p} \frac{|\psi(x) - \psi(y)|}{|x - y|^{N+sp}},$$

thanks to Young inequality and to the 1-Lipschitz character of f_ε .

By using this estimate, up to now we gained

$$\left(\int_{\Omega} (u_M + \delta)^{\beta \frac{N}{N-sp}} dx \right)^{\frac{N-sp}{N}} \leq C \frac{\|F\|_{L^q(\Omega)}}{\beta} \left(\frac{\beta + p - 1}{p \delta^{\frac{p-1}{p}}} \right)^p \left(\int_{\Omega} (u_M + \delta)^{\beta q'} dx \right)^{\frac{1}{q'}} + C \delta^{\beta} |\Omega|^{\frac{N-sp}{N}},$$

possibly for a different constant $C = C(N, p, s) > 0$. On the other hand, it is easy to see that³

$$\delta^{\beta} |\Omega|^{\frac{N-sp}{N}} \leq \frac{1}{\beta} \left(\frac{\beta + p - 1}{p} \right)^p |\Omega|^{1 - \frac{1}{q'} - \frac{sp}{N}} \left(\int_{\Omega} (u_M + \delta)^{\beta q'} dx \right)^{\frac{1}{q'}}.$$

Thus we get

$$(3.5) \quad \left(\int_{\Omega} (u_M + \delta)^{\beta \frac{N}{N-sp}} dx \right)^{\frac{N-sp}{N}} \leq C \frac{1}{\beta} \left(\frac{\beta + p - 1}{p} \right)^p \left(\int_{\Omega} (u_M + \delta)^{\beta q'} dx \right)^{\frac{1}{q'}} \times \left[\frac{\|F\|_{L^q(\Omega)}}{\delta^{p-1}} + |\Omega|^{1 - \frac{1}{q'} - \frac{sp}{N}} \right],$$

with $C = C(N, s, p) > 0$ as usual. We then choose $\delta > 0$ by

$$(3.6) \quad \delta = \|F\|_{L^q(\Omega)}^{\frac{1}{p-1}} |\Omega|^{-\frac{1}{p-1} (1 - \frac{1}{q'} - \frac{sp}{N})},$$

and set $\vartheta = \beta q'$. The previous inequality can be written as

$$\left(\int_{\Omega} (u_M + \delta)^{\chi \vartheta} dx \right)^{\frac{1}{\chi \vartheta}} \leq \left[C |\Omega|^{1 - \frac{1}{q'} - \frac{sp}{N}} \right]^{\frac{q'}{\vartheta}} \left(\frac{q'}{\vartheta} \right)^{\frac{q'}{\vartheta}} \left(\frac{\vartheta + q'p - q'}{q'p} \right)^{\frac{p q'}{\vartheta}} \left(\int_{\Omega} (u_M + \delta)^{\vartheta} dx \right)^{\frac{1}{\vartheta}},$$

where for simplicity

$$(3.7) \quad \chi = \frac{N}{N - sp} \frac{1}{q'} > 1.$$

We now iterate the previous inequality, by taking the following sequence of exponents

$$\vartheta_0 = 1, \quad \vartheta_{n+1} = \chi \vartheta_n = \chi^{n+1}.$$

Since $\chi > 1$, then

$$\sum_{n=0}^{\infty} \frac{1}{\vartheta_n} = \sum_{n=0}^{\infty} \frac{1}{\chi^n} = \frac{\chi}{\chi - 1} = \frac{N}{N - N q' + sp q'},$$

and

$$\prod_{n=0}^{\infty} \left(\frac{q'}{\vartheta_n} \right)^{\frac{q'}{\vartheta_n}} \left(\frac{\vartheta_n + q'p - q'}{q'p} \right)^{\frac{p q'}{\vartheta_n}} < +\infty.$$

By starting from $n = 0$, at the step n we have

$$\|u_M + \delta\|_{L^{\vartheta_{n+1}}} \leq \left[C |\Omega|^{1 - \frac{1}{q'} - \frac{sp}{N}} \right]^{\sum_{i=0}^n \frac{q'}{\vartheta_i}} \prod_{i=0}^n \left(\frac{q'}{\vartheta_i} \right)^{\frac{q'}{\vartheta_i}} \left(\frac{\vartheta_i + q'p - q'}{q'p} \right)^{\frac{p q'}{\vartheta_i}} \|u_M + \delta\|_{L^1}.$$

By taking the limit as n goes to ∞ we finally obtain

$$\|u_M\|_{L^\infty} \leq C \left(|\Omega|^{1 - \frac{1}{q'} - \frac{sp}{N}} \right)^{q' \frac{\chi}{\chi - 1}} \|u_M + \delta\|_{L^1},$$

³We use that

$$\frac{1}{\beta} \left(\frac{\beta + p - 1}{p} \right)^p \geq 1,$$

see Lemma A.5.

for some constant $C = C(N, p, s, q) > 0$. In particular, since $u_M \leq |u|$ and by triangle inequality, we get

$$\|u_M\|_{L^\infty} \leq C \left(|\Omega|^{1-\frac{1}{q'}-\frac{sp}{N}} \right)^{q' \frac{\chi}{\chi-1}} \left[\|u\|_{L^1} + \delta |\Omega| \right].$$

By recalling the definitions (3.6) and (3.7), we finally get the desired conclusion by letting M go to ∞ . \square

Remark 3.2. For eigenfunctions, starting from (3.4) one can reproduce the proof in [4, Theorem 3.1] for the first eigenfunction and prove that $u \in L^\infty(\Omega)$. This gives another proof of [14, Theorem 3.2], this time based on Moser's iterations. For $sp < N$ there holds

$$\|u\|_{L^\infty(\Omega)} \leq C'_{N,p,s} \lambda^{\frac{N}{sp^2}} \|u\|_{L^p(\Omega)}.$$

By Lebesgue interpolation, the previous also gives

$$\|u\|_{L^\infty(\Omega)} \leq (C'_{N,p,s})^p \lambda^{\frac{N}{sp}} \|u\|_{L^1(\Omega)}.$$

3.2. Caccioppoli inequalities. The previous result is based on the fact that if u is a solution of (3.2), then $|u|$ is a subsolution of the same equation. Like in the local case, this is a general fact which remains true for every $f(u)$ with f convex. This is the content of the next results.

Lemma 3.3 (Subsolutions, part I). *Let $1 < p < \infty$ and $0 < s < 1$. Let $F \in L^q(\Omega)$ for $q > N/(sp)$ and let $u \in \widetilde{W}_0^{s,p}(\Omega) \cap L^\infty(\Omega)$ be the solution of (3.2). Then for every $f : \mathbb{R} \rightarrow \mathbb{R}$ convex C^1 , the composition $v = f \circ u$ verifies*

$$(3.8) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy \leq \int_{\Omega} F |f'(u)|^{p-2} f'(u) \varphi dx,$$

for every positive function $\varphi \in \widetilde{W}_0^{s,p}(\Omega)$.

Proof. Observe that the assumption $u \in L^\infty(\Omega)$ is not restrictive, by Theorem 3.1. The proof goes exactly like in the first part of Theorem 3.1. We insert in (3.2) the test function $\varphi = \psi |f'(u)|^{p-2} f'(u)$, where $\psi \in C_0^\infty(\Omega)$ is a positive function. If f is not regular enough, a standard smoothing argument will be needed, we leave the details to the reader. Then by appealing to (A.1), we get (3.8) for smooth test functions. A density argument gives again the conclusion. \square

The following is very similar, but we lower the hypothesis on F .

Lemma 3.4 (Subsolutions, part II). *Let $1 < p < \infty$ and $0 < s < 1$ be such that $sp < N$. Let $F \in L^{(p^*)}'(\Omega)$ and $u \in \widetilde{W}_0^{s,p}(\Omega)$ be the solution of (3.2). Then for every $f : \mathbb{R} \rightarrow \mathbb{R}$ convex and L -Lipschitz, the composition $v = f \circ u$ verifies*

$$(3.9) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy \leq L^{p-1} \int_{\Omega} |F| \varphi dx,$$

for every positive function $\varphi \in \widetilde{W}_0^{s,p}(\Omega)$.

The following can be seen as the Moser-type counterpart of [8, Theorem 1.4], for equations with a right-hand side F .

Proposition 3.5 (Localized Caccioppoli inequality). *Let $1 < p < \infty$ and $0 < s < 1$ be such that $sp < N$ and let $F \in L^{(p^*)}'(\Omega)$. We assume that $v = f(u)$ is a subsolution, i.e. a function satisfying (3.9) for an L -Lipschitz convex function f and such that*

$$(3.10) \quad v \geq 0 \quad \text{on } \Omega' \Subset \Omega.$$

For every $\beta \geq 1$ and $\delta \geq 0$, we take

$$g(t) = (t + \delta)^\beta \quad \text{and} \quad G(t) = \int_0^t g'(\tau)^{\frac{1}{p}} d\tau = \frac{p \beta^{\frac{1}{p}}}{\beta + p - 1} (t + \delta)^{\frac{\beta + p - 1}{p}}, \quad t \geq 0.$$

Then for every positive $\psi \in C_0^\infty(\Omega)$ such that $\text{spt}(\psi) \Subset \Omega'$ we have

$$\begin{aligned} (3.11) \quad & \int_{\Omega'} \int_{\Omega'} \frac{|G(v(x)) \psi(x) - G(v(y)) \psi(y)|^p}{|x - y|^{N+sp}} dx dy \\ & \leq \frac{C}{\beta} \left(\frac{\beta + p - 1}{p} \right)^p \int_{\Omega'} \int_{\Omega'} \frac{|\psi(x) - \psi(y)|^p}{|x - y|^{N+sp}} (G(v(x))^p + G(v(y))^p) dx dy \\ & + C \left(\sup_{y \in \text{spt}(\psi)} \int_{\mathbb{R}^N \setminus \Omega'} \frac{|v(x)|^{p-1}}{|x - y|^{N+sp}} dx \right) \int_{\Omega'} g(v) \psi^p dx \\ & + C L^{p-1} \int_{\Omega} |F| g(v) \psi^p dx, \end{aligned}$$

for some constant $C = C(p) > 0$.

Proof. We insert in (3.9) the test function $\varphi = \psi^p g(v)$, where $\psi \in C_0^\infty(\Omega)$ is a positive function such that $\text{spt}(\psi) \Subset \Omega'$. Then we get

$$\begin{aligned} & \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} (g(v(x)) \psi(x)^p - g(v(y)) \psi(y)^p) dx dy \\ & \leq \int_{\Omega} |F| |f'(u)|^{p-1} g(v) \psi^p dx. \end{aligned}$$

We now split the double integral in three parts:

$$\begin{aligned} \mathcal{I}_1 &= \int_{\Omega'} \int_{\Omega'} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} (g(v(x)) \psi(x)^p - g(v(y)) \psi(y)^p) dx dy, \\ \mathcal{I}_2 &= \int_{\mathbb{R}^N \setminus \Omega'} \int_{\Omega'} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} g(v(x)) \psi(x)^p dx dy, \end{aligned}$$

and

$$\mathcal{I}_3 = - \int_{\Omega'} \int_{\mathbb{R}^N \setminus \Omega'} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} g(v(y)) \psi(y)^p dx dy$$

Estimate of \mathcal{I}_1 . For the first integral \mathcal{I}_1 , we proceed as follows: at first, for every $x, y \in \Omega'$ we have

$$\begin{aligned} & |v(x) - v(y)|^{p-2} (v(x) - v(y)) (g(v(x)) \psi(x)^p - g(v(y)) \psi(y)^p) \\ & = |v(x) - v(y)|^{p-2} (v(x) - v(y)) \frac{g(v(x)) - g(v(y))}{2} (\psi(x)^p + \psi(y)^p) \\ & + |v(x) - v(y)|^{p-2} (v(x) - v(y)) \frac{g(v(x)) + g(v(y))}{2} (\psi(x)^p - \psi(y)^p). \\ & \geq |v(x) - v(y)|^{p-2} (v(x) - v(y)) \frac{g(v(x)) - g(v(y))}{2} (\psi(x)^p + \psi(y)^p) \\ & - |v(x) - v(y)|^{p-1} \frac{g(v(y)) + g(v(x))}{2} |\psi(x)^p - \psi(y)^p|. \end{aligned}$$

In order to estimate the last term, we use at first

$$\begin{aligned}
 (3.12) \quad |\psi(x)^p - \psi(y)^p| &\leq p \max \{ \psi(x)^{p-1}, \psi(y)^{p-1} \} |\psi(x) - \psi(y)| \\
 &\leq p \left(\psi(x)^p + \psi(y)^p \right)^{\frac{p-1}{p}} |\psi(x) - \psi(y)|.
 \end{aligned}$$

Then we appeal to the definition of $g(t)$ and to Young inequality with exponents p and p' : thus we have

$$\begin{aligned}
 |v(x) - v(y)|^{p-1} \frac{(v(y) + \delta)^\beta + (v(x) + \delta)^\beta}{2} |\psi(x)^p - \psi(y)^p| \\
 \leq \frac{p}{2} |v(x) - v(y)|^{p-1} (v(x) + \delta)^\beta (\psi(x)^p + \psi(y)^p)^{\frac{p-1}{p}} |\psi(x) - \psi(y)| \\
 + \frac{p}{2} |v(x) - v(y)|^{p-1} (v(y) + \delta)^\beta (\psi(x)^p + \psi(y)^p)^{\frac{p-1}{p}} |\psi(x) - \psi(y)| \\
 \leq \varepsilon \frac{p-1}{2} |v(x) - v(y)|^p (v(x) + \delta)^{\beta-1} (\psi(x)^p + \psi(y)^p) \\
 + \frac{\varepsilon^{-\frac{1}{p-1}}}{2} (v(x) + \delta)^{\beta+p-1} |\psi(x) - \psi(y)|^p \\
 + \varepsilon \frac{p-1}{2} |v(x) - v(y)|^p (v(y) + \delta)^{\beta-1} (\psi(x)^p + \psi(y)^p) \\
 + \frac{\varepsilon^{-\frac{1}{p-1}}}{2} (v(y) + \delta)^{\beta+p-1} |\psi(x) - \psi(y)|^p,
 \end{aligned}$$

where $\varepsilon > 0$ will be chosen in a while. For the moment we have shown

$$\begin{aligned}
 |v(x) - v(y)|^{p-2} (v(x) - v(y)) (g(v(x)) \psi(x)^p - g(v(y)) \psi(y)^p) \\
 \geq |v(x) - v(y)|^{p-2} (v(x) - v(y)) \frac{g(v(x)) - g(v(y))}{2} (\psi(x)^p + \psi(y)^p) \\
 - \varepsilon \frac{p-1}{2} |v(x) - v(y)|^p \left((v(x) + \delta)^{\beta-1} + (v(y) + \delta)^{\beta-1} \right) (\psi(x)^p + \psi(y)^p) \\
 - \frac{\varepsilon^{-\frac{1}{p-1}}}{2} \left((v(x) + \delta)^{\beta+p-1} + (v(y) + \delta)^{\beta+p-1} \right) |\psi(x) - \psi(y)|^p.
 \end{aligned}$$

Now we can use the pointwise inequality (A.4) for the second term in the right-hand side, with the choices

$$a = v(x) + \delta \quad \text{and} \quad b = v(y) + \delta.$$

This gives us⁴

$$\begin{aligned}
 |v(x) - v(y)|^{p-2} (v(x) - v(y)) (g(v(x)) \psi(x)^p - g(v(y)) \psi(y)^p) \\
 \geq \frac{1 - 2\varepsilon(p-1)}{2} |v(x) - v(y)|^{p-2} (v(x) - v(y)) (g(v(x)) - g(v(y))) (\psi(x)^p + \psi(y)^p) \\
 - \frac{\varepsilon^{-\frac{1}{p-1}}}{2} \left((v(x) + \delta)^{\beta+p-1} + (v(y) + \delta)^{\beta+p-1} \right) |\psi(x) - \psi(y)|^p.
 \end{aligned}$$

⁴We use that the constant $\max\{1, 3 - \beta\}$ appearing in (A.4) is less than 2.

If we now choose $\varepsilon = (4(p-1))^{-1}$, use (A.3) and recall the definition of G , we finally get

$$\begin{aligned} & |v(x) - v(y)|^{p-2} (v(x) - v(y)) (g(v(x)) \psi(x)^p - g(v(y)) \psi(y)^p) \\ & \geq c |G(v(x)) - G(v(y))|^p (\psi(x)^p + \psi(y)^p) \\ & \quad - \frac{C}{\beta} \left(\frac{\beta + p - 1}{p} \right)^p (G(v(x))^p + G(v(y))^p) |\psi(x) - \psi(y)|^p, \end{aligned}$$

for some $c = c(p) > 0$ and $C = C(p) > 0$. If we further use

$$(3.13) \quad |G(v(x)) - G(v(y))|^p (\psi(x)^p + \psi(y)^p) \geq 2^{1-p} \left| G(v(x)) \psi(x) - G(v(y)) \psi(y) \right|^p,$$

this gives

$$\begin{aligned} (3.14) \quad & c \int_{\Omega'} \int_{\Omega'} \frac{\left| G(v(x)) \psi(x) - G(v(y)) \psi(y) \right|^p}{|x - y|^{N+sp}} dx dy \leq \mathcal{I}_1 \\ & + \frac{C}{\beta} \left(\frac{\beta + p - 1}{p} \right)^p \int_{\Omega'} \int_{\Omega'} \frac{|\psi(x) - \psi(y)|^p}{|x - y|^{N+sp}} (G(v(x))^p + G(v(y))^p) dx dy. \end{aligned}$$

Estimate of \mathcal{I}_2 . This is easier to estimate, we simply observe that by monotonicity of $\tau \mapsto |\tau|^{p-2} \tau$ and hypothesis (3.10), for $x \in \Omega'$ we have

$$|v(x) - v(y)|^{p-2} (v(x) - v(y)) \geq -|v(y)|^{p-2} v(y),$$

thus we get

$$\begin{aligned} (3.15) \quad \mathcal{I}_2 & \geq - \int_{\mathbb{R}^N \setminus \Omega'} \int_{\Omega'} \frac{|v(y)|^{p-2} v(y)}{|x - y|^{N+sp}} g(v(x)) \psi(x)^p dx dy \\ & \geq - \left(\sup_{x \in \text{spt}(\psi)} \int_{\mathbb{R}^N \setminus \Omega'} \frac{|v(y)|^{p-1}}{|x - y|^{N+sp}} dy \right) \int_{\Omega'} g(v(x)) \psi(x)^p dx. \end{aligned}$$

Estimate of \mathcal{I}_3 . This is estimated exactly in the same way as before, i.e. it is sufficient to use for $y \in \Omega'$

$$|v(x) - v(y)|^{p-2} (v(x) - v(y)) \leq |v(x)|^{p-2} v(x),$$

then

$$\begin{aligned} (3.16) \quad \mathcal{I}_3 & \geq - \int_{\Omega'} \int_{\mathbb{R}^N \setminus \Omega'} \frac{|v(x)|^{p-2} v(x)}{|x - y|^{N+sp}} g(v(y)) \psi(y)^p dx dy \\ & \geq - \left(\sup_{y \in \text{spt}(\psi)} \int_{\mathbb{R}^N \setminus \Omega'} \frac{|v(x)|^{p-1}}{|x - y|^{N+sp}} dx \right) \int_{\Omega'} g(v(y)) \psi(y)^p dy. \end{aligned}$$

Conclusion. Since we have

$$\sum_{i=1}^3 \mathcal{I}_i \leq L^{p-1} \int_{\Omega} |F| g(v) \psi^p dx,$$

it is sufficient to use (3.14), (3.15) and (3.16) in order to get the desired conclusion. \square

As in [8], we define the *nonlocal tail* of a function φ by

$$\text{Tail}(\varphi; x_0, R) := \left(R^{sp} \int_{\mathbb{R}^N \setminus B_R(x_0)} \frac{|\varphi(x)|^{p-1}}{|x - x_0|^{N+sp}} dx \right)^{\frac{1}{p-1}}.$$

Observe that this is a scaling invariant quantity. An important consequence of the Caccioppoli inequality is the following result for solutions (see [20, Lemma 2.2] for the case $F \equiv 0$). We state the result for the subconformal case $sp < N$, then it will be evident how to adapt the argument to cover the case $sp = N$, where $F \in L^q(\Omega)$ with $q > 1$.

Corollary 3.6. *Let $1 < p < \infty$ and $0 < s < 1$ such that $sp < N$ and let $F \in L^{(p^*)'}(\Omega)$. We take $0 < r < R$ such that $B_r(x_0) \subseteq B_R(x_0) \subseteq \Omega$. For the solution u of (3.2) there holds*

$$(3.17) \quad \begin{aligned} & \int_{B_r} \int_{B_r} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \\ & \leq \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^{N+sp+p} \left\{ \int_{B_R} |u|^p dx + R^N [\text{Tail}(u; x_0, R)]^p \right\} + C \|F\|_{L^{(p^*)'}(B_R)}^{\frac{p}{p-1}}, \end{aligned}$$

for some constant $C = C(N, p, s) > 0$.

Proof. In (3.11) we choose $\Omega' = B_R$ and $\psi \in C_0^\infty(B_{(r+R)/2})$ such that

$$0 \leq \psi \leq 1 \quad \text{in } B_{(r+R)/2}, \quad \psi \equiv 1 \text{ in } B_r \quad \text{and} \quad |\nabla \psi| \leq \frac{C}{R-r},$$

for some universal constant $C > 0$. We also take $v = u_+$ the positive part of u and $g(t) = t$, i.e. $\beta = 1$ and $\delta = 0$. Then from (3.11) we get

$$(3.18) \quad \begin{aligned} & \int_{B_R} \int_{B_R} \frac{|u_+(x)\psi(x) - u_+(y)\psi(y)|^p}{|x - y|^{N+sp}} dx dy + \frac{1}{R^{sp}} \int_{B_R} u_+^p \psi^p dx \\ & \leq \frac{C}{(R-r)^p} \int_{B_R} \int_{B_R} \frac{1}{|x - y|^{N+sp-p}} (u_+(x)^p + u_+(y)^p) dx dy + \frac{1}{R^{sp}} \int_{B_R} u_+^p \psi^p dx \\ & \quad + C \left(\sup_{y \in B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{u_+^{p-1}}{|x - y|^{N+sp}} dx \right) \int_{B_R} u_+ \psi^p dx + \int_{\Omega} |F| u_+ \psi^p dx. \end{aligned}$$

where we added the term

$$\frac{1}{R^{sp}} \int_{B_R} u_+^p \psi^p dx,$$

on both sides. We then observe that

$$(3.19) \quad \begin{aligned} & \int_{B_R} \int_{B_R} \frac{1}{|x - y|^{N+sp-p}} (u_+(x)^p + u_+(y)^p) dx dy \\ & = 2 \int_{B_R} \int_{B_R} \frac{u_+(x)^p}{|x - y|^{N+sp-p}} dx dy \\ & \leq 2 \int_{B_R} \left(\int_{B_{2R}} \frac{dy}{|x_0 - y|^{N+sp-p}} \right) u_+^p dx = c R^{p(1-s)} \int_{B_R} u_+^p dx, \end{aligned}$$

for some constant $c = c(N, s, p) > 0$. For the second term on the right-hand side of (3.18), we have

$$(3.20) \quad \begin{aligned} \sup_{y \in B_{(R+r)/2}} \int_{\mathbb{R}^N \setminus B_R} \frac{u_+^{p-1}}{|x - y|^{N+sp}} dx & \leq \frac{2^{N+sp} R^{N+sp}}{(R-r)^{N+sp}} \int_{\mathbb{R}^N \setminus B_R} \frac{u_+^{p-1}}{|x - x_0|^{N+sp}} dx \\ & = \frac{2^{N+sp} R^N}{(R-r)^{N+sp}} [\text{Tail}(u_+; x_0, R)]^{p-1}, \end{aligned}$$

since for every $y \in B_{(R+r)/2}$ and $x \in \mathbb{R}^N \setminus B_R$ we have

$$\begin{aligned} |x - y| &\geq |x - x_0| - |x_0 - y| \geq |x - x_0| - \frac{R+r}{2} \\ &\geq |x - x_0| - \frac{R+r}{2R} |x - x_0| = \frac{R-r}{2R} |x - x_0|. \end{aligned}$$

Finally, by Hölder and Young inequalities

$$\int_{\Omega} |F| u_+ \psi^p dx \leq \frac{(p-1)}{p} \frac{1}{\tau^{\frac{1}{p-1}}} \left(\int_{B_R} (|F| \psi^{p-1})^{(p^*)'} dx \right)^{\frac{p'}{(p^*)'}} + \frac{\tau}{p} \left(\int_{B_R} (u_+ \psi)^{p^*} dx \right)^{\frac{p}{p^*}}.$$

For the last term we use Sobolev inequality (2.6) and choose $\tau \ll 1$ small enough, in order to absorb it in the left-hand side of (3.18). Then from (3.18), (3.19) and (3.20) we get

$$\begin{aligned} &\int_{B_R} \int_{B_R} \frac{|u_+(x) \psi(x) - u_+(y) \psi(y)|^p}{|x - y|^{N+sp}} dx dy + \frac{1}{R^{sp}} \int_{B_R} u_+^p \psi^p dx \\ &\leq \frac{C}{R^{sp}} \left[\left(\frac{R}{R-r} \right)^p + 1 \right] \int_{B_R} u_+^p dx dy \\ &\quad + \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^{N+sp} \left[\text{Tail}(u_+; x_0, R) \right]^{p-1} \int_{B_R} u_+ dx + C \|F\|_{L^{(p^*)'}(B_R)}^{\frac{p}{p-1}}. \end{aligned}$$

By Young and Hölder inequalities we have

$$\left[\text{Tail}(u_+; x_0, R) \right]^{p-1} \int_{B_R} u_+ dx \leq \frac{(p-1) R^N}{p} \left[\text{Tail}(u_+; x_0, R) \right]^p + \frac{C}{p} \int_{B_R} u_+^p dx.$$

By using that $\psi \equiv 1$ on B_r and also $R/(R-r) > 1$ we thus get

$$\begin{aligned} &\int_{B_r} \int_{B_r} \frac{|u_+(x) - u_+(y)|^p}{|x - y|^{N+sp}} dx dy \\ &\leq \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^{N+sp+p} \left\{ \int_{B_R} u_+^p dx + R^N \left[\text{Tail}(u_+; x_0, R) \right]^p \right\} + C \|F\|_{L^{(p^*)'}(B_R)}^{\frac{p}{p-1}}, \end{aligned}$$

for some constant $C = C(N, p, s) > 0$. This proves (3.17). By reproducing the proof above for $v = u_-$, the negative part of u , and then summing up the resulting inequalities, we finally get inequality (3.17) for the solution itself. \square

Remark 3.7. We observe that the previous estimate still holds for $u - c$, where $c \in \mathbb{R}$. In particular, if we take $c = \bar{u}_{x_0, R}$, we get

$$\begin{aligned} (3.21) \quad &\int_{B_r} \int_{B_r} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \leq \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^{N+sp+p} \\ &\quad \times \left\{ \int_{B_R} |u - \bar{u}_{x_0, R}|^p dx + R^N \left[\text{Tail}(u - \bar{u}_{x_0, R}; x_0, R) \right]^p \right\} + C \|F\|_{L^{(p^*)'}(B_R)}^{\frac{p}{p-1}}. \end{aligned}$$

3.3. Local L^∞ estimates. In the homogeneous case, the local L^∞ estimate below has been proved in [8, Theorem 1.1]. The proof uses a De Giorgi-type iteration. The case $F \not\equiv 0$ has been treated in [20, Theorem 1.2], by using a perturbative argument. The result in [20] provides a pointwise nonlinear potential estimate on the solution.

Our approach is more elementary and direct, it relies on the Caccioppoli inequality of Proposition 3.5 and a Moser's iteration.

Theorem 3.8 (Local boundedness). *Let $1 < p < \infty$ and $0 < s < 1$ such that $sp \leq N$. Let $F \in L^q(\Omega)$ for $q > N/(sp)$ and $v = f(u)$ be a subsolution, i.e. a function satisfying (3.9) with f convex and L -Lipschitz. We take $0 < r_0 < R_0$ such that $B_{r_0}(x_0) \subseteq B_{R_0}(x_0) \subseteq \Omega$, and suppose that*

$$v \geq 0 \quad \text{on } B_{R_0}(x_0).$$

Then the following scaling invariant estimate holds

$$(3.22) \quad \begin{aligned} \|v\|_{L^\infty(B_{R_0})} &\leq C \left(\frac{1}{|B_{R_0}|} \int_{B_{R_0}} v^p dx \right)^{\frac{1}{p}} + C \left(\frac{R_0}{r_0} \right)^{\frac{sp}{p-1}} \text{Tail}(v; x_0, r_0) \\ &\quad + C L \left(R_0^{sp - \frac{N}{q}} \|F\|_{L^q(B_{R_0})} \right)^{\frac{1}{p-1}}, \end{aligned}$$

where

$$C = C_0 \left(\frac{R_0}{r_0} \right)^{\frac{N}{p}} \left(\frac{R_0}{R_0 - r_0} \right)^{(N+sp+p) \frac{N}{sp^2}} \quad \text{and} \quad C_0 = C_0(N, s, p, q) > 0.$$

Proof. We use Proposition 3.5 in order to perform a suitable Moser's iteration. We first prove (3.22) for $r_0 = 1$ and $R_0 = \gamma > 1$, then we will get (3.22) for general $r_0 < R_0$ by a scaling argument. It is also clear that it is sufficient to prove the result for $L = 1$, up to replace v by v/L .

Estimate at scale 1. We thus take $1 = r_0 \leq r < R \leq \gamma$, choose $\Omega' = B_R$ and $\psi \in C_0^\infty(B_{(r+R)/2})$ such that

$$0 \leq \psi \leq 1 \quad \text{in } B_{(r+R)/2}, \quad \psi \equiv 1 \text{ in } B_r \quad \text{and} \quad |\nabla \psi| \leq \frac{C}{R-r},$$

for some universal constant $C > 0$. For notational simplicity, we also set

$$v_\delta = v + \delta.$$

From (3.11) we get

$$(3.23) \quad \begin{aligned} &\beta \left(\frac{p}{\beta + p - 1} \right)^p \int_{B_R} \int_{B_R} \frac{|v_\delta(x)^{\frac{\beta+p-1}{p}} \psi(x) - v_\delta(y)^{\frac{\beta+p-1}{p}} \psi(y)|^p}{|x-y|^{N+sp}} dx dy \\ &\leq \frac{C}{(R-r)^p} \int_{B_R} \int_{B_R} \frac{1}{|x-y|^{N+sp-p}} \left(v_\delta(x)^{\beta+p-1} + v_\delta(y)^{\beta+p-1} \right) dx dy \\ &\quad + C \left(\sup_{y \in B_{\frac{R+r}{2}}} \int_{\mathbb{R}^N \setminus B_R} \frac{|v(x)|^{p-1}}{|x-y|^{N+sp}} dx \right) \int_{B_R} v_\delta^\beta \psi^p dx + \int_\Omega |F| v_\delta^\beta \psi^p dx. \end{aligned}$$

We then observe that by proceeding as in (3.19), we have

$$(3.24) \quad \int_{B_R} \int_{B_R} \frac{(v_\delta(x)^{\beta+p-1} + v_\delta(y)^{\beta+p-1})}{|x-y|^{N+sp-p}} dx dy \leq c R^{p(1-s)} \int_{B_R} v_\delta^{\beta+p-1} dx,$$

for some constant $c = c(N, s, p) > 0$. As for the second term in the right-hand side of (3.23), as in (3.20) we have (recall that $r_0 = 1$)

$$(3.25) \quad \sup_{y \in B_{(R+r)/2}} \int_{\mathbb{R}^N \setminus B_R} \frac{|v(x)|^{p-1}}{|x-y|^{N+sp}} dx \leq \frac{2^{N+sp} R^{N+sp}}{(R-r)^{N+sp}} [\text{Tail}(v; x_0, 1)]^{p-1}.$$

By collecting (3.24) and (3.25) in (3.23), we get

$$\begin{aligned} & \beta \left(\frac{p}{\beta + p - 1} \right)^p \int_{B_R} \int_{B_R} \frac{\left| v_\delta(x)^{\frac{\beta+p-1}{p}} \psi(x) - v_\delta(y)^{\frac{\beta+p-1}{p}} \psi(y) \right|^p}{|x-y|^{N+sp}} dx dy \\ & \leq \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^p \int_{B_R} v_\delta^{\beta+p-1} dx \\ & \quad + C \left(\frac{R}{R-r} \right)^{N+sp} \left[\text{Tail}(v; x_0, 1) \right]^{p-1} \int_{B_R} v_\delta^\beta \psi^p dx + \int_{B_R} |F| v_\delta^\beta \psi^p dx. \end{aligned}$$

If we add the term

$$\beta \left(\frac{p}{\beta + p - 1} \right)^p \frac{1}{R^{sp}} \int_{B_R} v_\delta^{\beta+p-1} \psi^p dx,$$

on both sides and use again the Sobolev inequality (2.6), we gain⁵

$$\begin{aligned} & \beta \left(\frac{p}{\beta + p - 1} \right)^p \left(\int_{B_R} \left(v_\delta^{\frac{\beta+p-1}{p}} \psi \right)^{p^*} dx \right)^{\frac{p}{p^*}} \\ (3.26) \quad & \leq \frac{C}{R^{sp}} \left[\left(\frac{R}{R-r} \right)^p + 1 \right] \int_{B_R} v_\delta^{\beta+p-1} dx + C \int_{\Omega} |F| v_\delta^\beta \psi^p dx \\ & \quad + C \left(\frac{R}{R-r} \right)^{N+sp} \left[\text{Tail}(v; x_0, 1) \right]^{p-1} \int_{B_R} v_\delta^\beta dx, \end{aligned}$$

for some constant $C = C(N, s, p) > 0$. For the term containing F in (3.26), we use

$$(3.27) \quad \delta^{p-1} v_\delta^\beta \leq v_\delta^{\beta+p-1},$$

Hölder inequality and Lebesgue interpolation⁶. These yield, for a parameter $0 < \tau \ll 1$

$$\begin{aligned} \int_{\Omega} |F| v_\delta^\beta \psi^p dx & \leq \delta^{1-p} \int_{\Omega} |F| v_\delta^{\beta+p-1} \psi^p dx \leq \delta^{1-p} \|F\|_{L^q(B_R)} \left(\int_{B_R} \left(v_\delta^{\frac{\beta+p-1}{p}} \psi \right)^{pq'} dx \right)^{\frac{1}{q'}} \\ & \leq c \tau \delta^{1-p} \|F\|_{L^q(B_R)} \left(\int_{B_R} \left(v_\delta^{\frac{\beta+p-1}{p}} \psi \right)^{p^*} dx \right)^{\frac{p}{p^*}} \\ & \quad + c \tau^{-\frac{N}{spq-N}} \delta^{1-p} \|F\|_{L^q(B_R)} \int_{B_R} v_\delta^{\beta+p-1} \psi^p dx, \end{aligned}$$

for $c = c(N, s, p, q) > 0$. By choosing τ as follows

$$\tau = \frac{\delta^{p-1}}{2c} \frac{1}{\|F\|_{L^q(B_R)}} \beta \left(\frac{p}{\beta + p - 1} \right)^p,$$

and observing that for $\beta \geq 1$, we have

$$\beta \left(\frac{p}{\beta + p - 1} \right)^p \geq \left(\frac{p}{\beta + p - 1} \right)^{p-1},$$

⁵We also use that

$$\beta \left(\frac{p}{\beta + p - 1} \right)^p \leq 1,$$

on the right-hand side, see Lemma A.5.

⁶Observe that we have $p < pq' < p^*$, thanks to the assumption on q .

from (3.26) and (3.27) we can infer

$$\begin{aligned} \left(\int_{B_r} v_\delta^\vartheta \frac{p^*}{p} dx \right)^{\frac{p}{p^*}} &\leq \left(\frac{\vartheta}{p} \right)^{p-1} \int_{B_R} v_\delta^\vartheta dx \\ &\times \left\{ \frac{C}{R^{sp}} \left[\left(\frac{R}{R-r} \right)^p + 1 \right] + \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^{N+sp} \left[\gamma^{\frac{sp}{p-1}} \frac{\text{Tail}(v; x_0, 1)}{\delta} \right]^{p-1} \right. \\ &\left. + \frac{C}{R^{sp}} \gamma^{sp} \left(\frac{\|F\|_{L^q(B_R)}}{\delta^{p-1}} \right)^{\frac{spq}{spq-N}} \left(\frac{\vartheta}{p} \right)^{(p-1) \frac{N}{spq-N}} \right\}, \end{aligned}$$

where we set $\vartheta = \beta + p - 1$. By observing that $R/(R-r) > 1$ and $\vartheta/p \geq 1$, from the previous we can also obtain

$$(3.28) \quad \left(\int_{B_r} v_\delta^\vartheta \frac{p^*}{p} dx \right)^{\frac{p}{p^*}} \leq \frac{C}{R^{sp}} \left(\frac{R}{R-r} \right)^{N+sp+p} \mathcal{T}(\delta) \left(\frac{\vartheta}{p} \right)^\eta \int_{B_R} v_\delta^\vartheta dx,$$

where $C = C(N, p, s, q) > 0$. In the previous estimate we set for simplicity

$$\eta = (p-1) \frac{spq}{spq-N} \quad \text{and} \quad \mathcal{T}(\delta) = 1 + \left[\gamma^{\frac{sp}{p-1}} \frac{\text{Tail}(v; x_0, 1)}{\delta} \right]^{p-1} + \left(\gamma^{\frac{sp}{\eta}} \frac{\|F\|_{L^q(B_R)}^{\frac{1}{p-1}}}{\delta} \right)^\eta.$$

If we define $\chi = p^*/p > 1$, then from (3.28) with simple manipulations we have

$$(3.29) \quad \|v_\delta\|_{L^{\vartheta\chi}(B_r)} \leq \left(C \left(\frac{R}{R-r} \right)^{N+sp+p} \mathcal{T}(\delta) \right)^{\frac{1}{\vartheta}} \left(\vartheta^{\frac{1}{\vartheta}} \right)^\eta \|v_\delta\|_{L^{\vartheta}(B_R)}.$$

We now define the sequences $\{\vartheta_k\}_{k \in \mathbb{N}}$ and $\{\varrho_k\}_{k \in \mathbb{N}}$ by

$$\vartheta_k = \chi \vartheta_{k-1} = \left(\frac{N}{N-sp} \right)^k p, \quad \varrho_k = 1 + \frac{\gamma-1}{2^k}.$$

Observe that

$$\left(\frac{\varrho_k}{\varrho_k - \varrho_{k+1}} \right)^{N+sp+p} \leq \left(2^{k+1} \right)^{N+sp+p} \left(\frac{\gamma}{\gamma-1} \right)^{N+sp+p},$$

thus (3.29) applied with $r = \varrho_{k+1}$ and $R = \varrho_k$ gives

$$(3.30) \quad \|v_\delta\|_{L^{\vartheta_{k+1}}(B_{\varrho_{k+1}})} \leq \left(C' \left(\frac{\gamma}{\gamma-1} \right)^{N+sp+p} \mathcal{T}(\delta) \right)^{\frac{1}{\vartheta_k}} \left(\vartheta_k^\eta 2^{k(N+(1+s)p)} \right)^{\frac{1}{\vartheta_k}} \|v_\delta\|_{L^{\vartheta_k}(B_{\varrho_k})},$$

where $C' = C'(N, s, p, q) > 0$. We now observe that

$$\sum_{k=0}^{\infty} \frac{1}{\vartheta_k} = \frac{1}{p} \sum_{k=0}^{\infty} \left(\frac{N-sp}{N} \right)^k = \frac{N}{sp^2},$$

and that

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(\prod_{k=0}^n \vartheta_k^{\frac{\eta}{\vartheta_k}} \left(2^{N+(1+s)p} \right)^{\frac{k}{\vartheta_k}} \right) &= \lim_{n \rightarrow \infty} \exp \left(\eta \sum_{k=0}^n \frac{\log \vartheta_k}{\vartheta_k} \right) \\ &\times \exp \left((N+(s+1)p) \log 2 \sum_{k=0}^n \frac{k}{\vartheta_k} \right) < +\infty, \end{aligned}$$

then by starting from $k = 0$ and iterating (3.30) infinitely many times, we finally get

$$\|v_\delta\|_{L^\infty(B_1)} \leq c' \left(\left(\frac{\gamma}{\gamma - 1} \right)^{N+sp+p} \mathcal{T}(\delta) \right)^{\frac{N}{sp^2}} \|v_\delta\|_{L^p(B_\gamma)}$$

for some constant $c' = c'(N, s, p, q) > 0$. If we now recall the definition of $\mathcal{T}(\delta)$, choose

$$\delta = \gamma^{\frac{sp}{p-1}} \text{Tail}(v; x_0, 1) + \|F\|_{L^q(B_\gamma)}^{\frac{1}{p-1}} \gamma^{\frac{sp}{q}},$$

and use the triangle inequality, from the previous we finally get (3.22) for $r_0 = 1$ and $R_0 = \gamma > 1$.

Estimate at general scales. We now take $r_0 < R_0$ and define

$$\gamma = \frac{R_0}{r_0} \quad \text{and} \quad \lambda = r_0.$$

The function $v_\lambda(x) := v(\lambda x)$ satisfies (3.8) with right-hand side $F_\lambda(x) := \lambda^{sp} F(\lambda x)$. It is now sufficient to use (3.22) for v_λ with balls B_1 and B_γ , together with the facts

$$\|v_\lambda\|_{L^\infty(B_1)} = \|v\|_{L^\infty(B_{r_0})}, \quad \int_{B_\gamma} v_\lambda^p dx = \left(\frac{R_0}{r_0} \right)^N \frac{1}{R_0^N} \int_{B_{R_0}} v^p dx,$$

$$\text{Tail}(v_\lambda; x_0, 1) = \text{Tail}(v; x_0, r_0) \quad \text{and} \quad \gamma^{sp - \frac{N}{q}} \|F_\lambda\|_{L^q(B_\gamma)} = R_0^{sp - \frac{N}{q}} \|F\|_{L^q(B_{R_0})}.$$

These give the desired conclusion. \square

3.4. Basic comparison estimates. We take $B_r := B_r(x_0) \Subset \Omega$ and consider the function $v \in W^{s,p}(\mathbb{R}^N)$ such that $v - u \in \widetilde{W}_0^{s,p}(B_r)$ and which solves the homogeneous equation

$$(3.31) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|v(x) - v(y)|^{p-2} (v(x) - v(y))}{|x - y|^{N+sp}} (\varphi(x) - \varphi(y)) dx dy = 0,$$

for every $\varphi \in \widetilde{W}_0^{s,p}(B_r)$. We point out that v is a locally Hölder function, see [8, Theorem 1.4]. It also satisfies a suitable Harnack inequality, as recently proved in [9]. We define

$$w = u - v \in \widetilde{W}_0^{s,p}(B_r),$$

then we have the following basic estimate. As usual, for simplicity we enunciate it only for the subconformal case $sp < N$, the reader should be able to adapt it to the conformal case $sp = N$, with $F \in L^q(\Omega)$ for some $q > 1$.

Lemma 3.9. *Let $1 < p < \infty$ and $0 < s < 1$ such that $sp < N$. For $F \in L^{(p^*)}'(\Omega)$ and every $B_{2r} := B_{2r}(x_0) \Subset \Omega$, with the notation above there holds*

$$(3.32) \quad \left(\int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{1}{p}} \leq \max\{2 - p, 0\} C \|F\|_{L^{(p^*)}'(B_r)} \left(\int_{B_{2r}} \int_{B_{2r}} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{2-p}{p}} + C \left(\|F\|_{L^{(p^*)}'(B_r)} \right)^{\frac{1}{p-1}},$$

for some constant $C = C(N, p, s) > 0$.

Proof. We will use for notational simplicity the function $J_p(t) = |t|^{p-2}t$. We test equations (3.2) and (3.31) with $\varphi = w$, subtracting them we get

$$(3.33) \quad \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{J_p(u(x) - u(y)) - J_p(v(x) - v(y))}{|x - y|^{N+sp}} (w(x) - w(y)) dx dy = \int_{\Omega} F w dx.$$

We now distinguish two cases.

Case $1 < p < 2$. We now observe that thanks to (B.5)

$$\begin{aligned} |w(x) - w(y)|^p &\leq c \left(J_p(u(x) - u(y)) - J_p(v(x) - v(y)) (w(x) - w(y)) \right)^{\frac{p}{2}} \\ &\quad \times \left((u(x) - u(y))^2 + (v(x) - v(y))^2 \right)^{\frac{2-p}{2} \frac{p}{2}}, \end{aligned}$$

thus by integrating this over B_{2r} , using Hölder inequality and (3.33) we get

$$\begin{aligned} &\int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \\ &\leq c \int_{B_{2r}} \int_{B_{2r}} \frac{\left(J_p(u(x) - u(y)) - J_p(v(x) - v(y)) (w(x) - w(y)) \right)^{\frac{p}{2}}}{|x - y|^{N+sp}} \\ &\quad \times \left((u(x) - u(y))^2 + (v(x) - v(y))^2 \right)^{\frac{2-p}{2} \frac{p}{2}} dx dy \\ &\leq c \left(\int_{B_{2r}} \int_{B_{2r}} \frac{J_p(u(x) - u(y)) - J_p(v(x) - v(y))}{|x - y|^{N+sp}} (w(x) - w(y)) dx dy \right)^{\frac{p}{2}} \\ &\quad \times \left(\int_{B_{2r}} \int_{B_{2r}} \frac{(|u(x) - u(y)|^2 + |v(x) - v(y)|^2)^{\frac{p}{2}}}{|x - y|^{N+sp}} dx dy \right)^{\frac{2-p}{2}} \\ &\leq c \left(\|F\|_{L^{(p^*)}'(B_r)} \|w\|_{L^{p^*}(B_r)} \right)^{\frac{p}{2}} \left(\int_{B_{2r}} \int_{B_{2r}} \frac{(|u(x) - u(y)|^2 + |v(x) - v(y)|^2)^{\frac{p}{2}}}{|x - y|^{N+sp}} dx dy \right)^{\frac{2-p}{2}}. \end{aligned}$$

Observe that in the right-hand side of (3.33) we used that $w \equiv 0$ outside B_r . Since $v = u - w$ and by subadditivity of $\tau \mapsto \tau^{p/2}$, we have

$$\begin{aligned} &\int_{B_{2r}} \int_{B_{2r}} \frac{(|u(x) - u(y)|^2 + |v(x) - v(y)|^2)^{\frac{p}{2}}}{|x - y|^{N+sp}} dx dy \\ &\leq C \int_{B_{2r}} \int_{B_{2r}} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy + C \int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy, \end{aligned}$$

for some constant $C = C(p) > 0$. Up to now, we obtained

$$\begin{aligned} &\int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \\ &\leq C \left(\|w\|_{L^{p^*}(B_r)} \right)^{\frac{p}{2}} \left(\|F\|_{L^{(p^*)}'(B_r)}^{\frac{p}{2-p}} \right)^{\frac{2-p}{2}} \\ &\quad \times \left(\int_{B_{2r}} \int_{B_{2r}} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy + \int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{2-p}{2}}, \end{aligned}$$

possibly for a different $C = C(p) > 0$. If we now use Young inequality with exponent 2 and the continuous embedding $\widetilde{W}_0^{s,p}(B_{2r}) \hookrightarrow L^{p^*}(B_r)$ of Proposition 2.3, from the previous and subadditivity of $\tau \mapsto \tau^{2-p}$ we get

$$\begin{aligned} & \int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \\ & \leq c \|F\|_{L^{(p^*)}'(B_r)}^p \left[\left(\int_{B_{2r}} \int_{B_{2r}} \frac{|u(x) - u(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{2-p} + \left(\int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{2-p} \right]. \end{aligned}$$

It is now sufficient to estimate the last term by means of Young inequality with exponents $1/(2-p)$ and $1/(p-1)$ in order to conclude.

Case $p \geq 2$. This case is simpler. In this case (B.6) implies

$$|w(x) - w(y)|^p \leq c \left(J_p(u(x) - u(y)) - J_p(v(x) - v(y)) \right) (w(x) - w(y)),$$

and thus from (3.33) we get

$$\begin{aligned} & \int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \leq \int_{B_r} F w dx \\ & \leq \|F\|_{L^{(p^*)}'(B_r)} \|w\|_{L^{p^*}(B_r)} \leq \frac{(p-1) \tau^{\frac{1}{1-p}}}{p} \|F\|_{L^{(p^*)}'(B_r)}^{\frac{p}{p-1}} + \frac{\tau}{p} \|w\|_{L^{p^*}(B_r)}^p. \end{aligned}$$

It is now sufficient to use again the Sobolev inequality of Proposition 2.3 to conclude. \square

Remark 3.10. The previous Lemma is the analogous of [20, Lemmas 3.2 & 3.4], under the simplified assumption that the right-hand side F belongs to $L^{(p^*)}'(\Omega)$. For $1 < p < 2$, the further restriction $p > 2 - s/N$ is needed in [20], since the right-hand side F is a measure not necessarily belonging to the relevant dual Sobolev space. On the contrary, under the standing assumption of this section we do not need this restriction on p .

We now introduce as in [20] the *global excess* functional

$$\mathcal{E}(u; x_0, r) = \left(\frac{1}{|B_r(x_0)|} \int_{B_r(x_0)} |u - \bar{u}_{x_0,r}|^p dx \right)^{\frac{1}{p}} + \text{Tail}(u - \bar{u}_{x_0,r}; x_0, r),$$

where $\bar{u}_{x_0,r}$ is the average of u over the ball $B_r(x_0)$, see (2.1). If we appeal to Corollary 3.6 and Remark 3.7, we get the following.

Lemma 3.11. *Let $1 < p < \infty$ and $0 < s < 1$ be such that $sp < N$ and let $F \in L^{(p^*)}'(\Omega)$. Then for every $B_{2r} := B_{2r}(x_0) \Subset B_R \Subset \Omega$, with the notation above there holds*

$$\begin{aligned} & \left(\int_{B_{2r}} \int_{B_{2r}} \frac{|w(x) - w(y)|^p}{|x - y|^{N+sp}} dx dy \right)^{\frac{1}{p}} \\ (3.34) \quad & \leq \max\{2-p, 0\} C \|F\|_{L^{(p^*)}'(B_R)} \left[R^{\frac{N}{p}-s} \left(\frac{R}{R-2r} \right)^{\frac{N}{p}+s+1} \right]^{2-p} \mathcal{E}(u; x_0, R)^{2-p} \\ & + C \left(\|F\|_{L^{(p^*)}'(B_R)} \right)^{\frac{1}{p-1}}, \end{aligned}$$

for some constant $C = C(N, p, s) > 0$. Moreover, we also have

$$(3.35) \quad \begin{aligned} \|w\|_{L^{p^*}(B_r)} &\leq \max\{2 - p, 0\} \tilde{C} \|F\|_{L^{(p^*)}'(B_R)} \left[R^{\frac{N}{p} - s} \left(\frac{R}{R - 2r} \right)^{\frac{N}{p} + s + 1} \right]^{2-p} \mathcal{E}(u; x_0, R)^{2-p} \\ &\quad + \tilde{C} \left(\|F\|_{L^{(p^*)}'(B_R)} \right)^{\frac{1}{p-1}}, \end{aligned}$$

where again $\tilde{C} = \tilde{C}(N, p, s) > 0$.

Proof. We notice at first that (3.35) follows from (3.34) and the Sobolev inequality of Proposition 2.3.

Let us prove (3.34). For $p \geq 2$ there is nothing to prove. For $1 < p < 2$, it is sufficient to combine (3.21) and (3.32), then with some simple computations one gets (3.34). \square

Remark 3.12. The previous Lemma is the analogue of [20, Lemmas 3.3 & 3.5], when F belongs to $L^{(p^*)}'(\Omega)$.

3.5. A theorem by Kuusi, Mingione and Sire.

Theorem 3.13 ([20], Theorem 1.5). *Let $1 < p < \infty$ and $0 < s < 1$. If $F \in L^q(\Omega)$ with $q > N/(sp)$, then the solution $u \in \widetilde{W}_0^{s,p}(\Omega)$ of (3.2) is continuous.*

Proof. As already explained, for $p \geq 2$ the continuity result of [20, Theorem 1.5] directly applies to our situation. For the case $1 < p < 2$, it is sufficient to reproduce the proof of [20, Theorem 1.5], by replacing their Lemmas 3.3 and 3.5 with our Lemma 3.11 above.

For the reader's convenience, we briefly explain the main points of the proof by Kuusi, Mingione and Sire, by referring to their paper for the details. The result in [20] is based on the following pointwise estimate⁷

$$(3.36) \quad |\bar{u}_{x_0, r} - u(x_0)| \leq c \left(r^{sq - \frac{N}{q}} \|F\|_{L^q(B_r)} \right)^{\frac{1}{p-1}} + c \mathcal{E}(u; x_0, r),$$

valid for every ball $B_r := B_r(x_0) \Subset \Omega$, see [20, Theorem 1.4]. Moreover, u enjoys the following VMO-type property, locally in Ω' in Ω : for every $\Omega'' \Subset \Omega'$

$$(3.37) \quad \lim_{\varrho_0 \rightarrow 0} \left[\sup_{x_0 \in \Omega''} \sup_{0 < \varrho < \varrho_0} \mathcal{E}(u; x_0, \varrho) \right] = 0.$$

Then (3.36) and (3.37) imply that u is the uniform limit of the net of continuous functions $\{\bar{u}_{x_0, r}\}_r$, thus u itself is continuous. In turn, both the proof (3.36) and that of (3.37) are based on the crucial

⁷In [20], the local term in the right-hand side is replaced by the *Wolff potential* $\mathbf{W}_{s,p}^F(x_0, r)$, which for a general Borel measure μ is defined by

$$\mathbf{W}_{s,p}^\mu(x_0, r) := \int_0^r \left(\frac{|\mu|(B_\varrho(x_0))}{\varrho^{N-sp}} \right)^{\frac{1}{p-1}} \frac{d\varrho}{\varrho}.$$

It is standard to see that when μ has a density $F \in L^q$ (here $q > N/(sp)$) with respect to the N -dimensional Lebesgue measure, we have

$$\mathbf{W}_{s,p}^\mu(x_0, r) \leq c \left(\|F\|_{L^q(B_r(x_0))} r^{sq - \frac{N}{q}} \right)^{\frac{1}{p-1}},$$

with $c = c(N, s, p, q) > 0$.

decay estimate

$$(3.38) \quad \begin{aligned} \mathcal{E}(u; x_0, \sigma r) &\leq c \sigma^\alpha \mathcal{E}(u; x_0, r) + c \left[\left(\frac{1}{\sigma} \right)^{\frac{N}{p-1}} + \left(\frac{1}{\sigma} \right)^{\frac{N}{p}} \right] \\ &\quad \times \left\{ \tau \mathcal{E}(u; x_0, 2r) + \tau^{\frac{p-2}{p-1}} \left(r^{s p - \frac{N}{q}} \|F\|_{L^q(B_{2r})} \right)^{\frac{1}{p-1}} \right\}, \end{aligned}$$

where $c = c(N, s, p) > 0$, $\alpha = \alpha(N, s, p) > 0$ is smaller than 1 and $0 < \sigma, \tau < 1$ are free parameters. We point out that this is equation (5.5) from [20]. The proof of (3.38) relies on a perturbative argument in order to transfer the decay estimate from v to u . As before, v is the solution of the homogeneous problem in the relevant ball, having u has boundary datum. It is precisely here that Lemma 3.11 for $w = u - v$ comes into play. \square

Corollary 3.14 (Eigenfunctions). *Let $1 < p < \infty$ and $0 < s < 1$. Every (s, p) -eigenfunction of the open bounded set $\Omega \subset \mathbb{R}^N$ is continuous.*

Proof. If $sp > N$ there is nothing to prove. For $sp \leq N$, we already know by Remark 3.2 that eigenfunctions are bounded. In particular an (s, p) -eigenfunction u solves (3.1) with $F = \lambda |u|^{p-2} u \in L^\infty(\Omega)$. Then the conclusion follows from Theorem 3.13. \square

4. THE SECOND EIGENVALUE

We start by defining

$$\lambda_2(\Omega) = \inf_{f \in \mathcal{C}_1(\Omega)} \max_{u \in \text{Im}(f)} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p,$$

where the set $\mathcal{C}_1(\Omega)$ is given by

$$\mathcal{C}_1(\Omega) = \{f : \mathbb{S}^1 \rightarrow \mathcal{S}_p(\Omega) : f \text{ odd and continuous}\}.$$

We have the following preliminary result.

Theorem 4.1. *The quantity $\lambda_2(\Omega)$ is an (s, p) -eigenvalue. Moreover we have $\lambda_1(\Omega) < \lambda_2(\Omega)$ and every eigenfunction $u \in \mathcal{S}_p(\Omega)$ associated to $\lambda_2(\Omega)$ has to change sign.*

Proof. We divide the proof in three parts.

Part 1: $\lambda_2(\Omega)$ is an eigenvalue. In order to prove that $\lambda_2(\Omega)$ is a critical point of the functional $\Phi_{s,p}$ defined in (2.8) on the manifold $\mathcal{S}_p(\Omega)$, it is sufficient to check that $\Phi_{s,p}$ verifies the Palais-Smale condition. The claim will then follow by applying [6, Proposition 2.7]. Let us take a sequence $\{u_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_p(\Omega)$ such that

$$(4.1) \quad \Phi_{s,p}(u_n) \leq C \quad \text{and} \quad \lim_{n \rightarrow \infty} \|D\Phi_{s,p}(u_n)|_{T_{u_n}\mathcal{S}_p(\Omega)}\|_* = 0,$$

where $D\Phi_{s,p}(u_n)$ denotes the differential of $\Phi_{s,p}$ at the point u_n and $T_{u_n}\mathcal{S}_p(\Omega)$ is the tangent space to $\mathcal{S}_p(\Omega)$ at the point u_n , given by

$$T_{u_n}\mathcal{S}_p(\Omega) = \left\{ \varphi \in \widetilde{W}_0^{s,p}(\Omega) : \int_{\Omega} |u_n|^{p-2} u_n \varphi \, dx = 0 \right\}.$$

Then the second hypothesis in (4.1) implies that there exists a sequence $\varepsilon_n > 0$ converging to 0 and such that

$$(4.2) \quad \left| D\Phi_{s,p}(u_n)[\varphi] \right| \leq \varepsilon_n \|\varphi\|_{\widetilde{W}_0^{s,p}(\Omega)}, \quad \text{for every } \varphi \in T_{u_n}\mathcal{S}_p(\Omega).$$

On the other hand, by the first hypothesis in (4.1) we can infer that $\{u_n\}_{n \in \mathbb{N}}$ is converging to a function u (up to a subsequence), strongly in $L^p(\Omega)$ and weakly in $\widetilde{W}_0^{s,p}(\Omega)$ (see for example [4,

Theorem 2.7]). By strong convergence in L^p we have of course $u \in \mathcal{S}_p(\Omega)$. Also observe that the sequence

$$\delta_n := \int_{\Omega} |u_n|^{p-2} u_n u \, dx,$$

converges to 1, as n goes to ∞ . We then define the new sequence $\{v_n\}_{n \in \mathbb{N}}$ by

$$v_n = \delta_n u_n - u, \quad n \in \mathbb{N},$$

and we observe that $v_n \in T_{u_n} \mathcal{S}_p(\Omega)$ for every n . Thus by (4.2) we get

$$\lim_{n \rightarrow \infty} \left| D\Phi_{s,p}(u_n)[u_n - u] \right| \leq \lim_{n \rightarrow \infty} \left| D\Phi_{s,p}(u_n)[v_n] \right| + \lim_{n \rightarrow \infty} |1 - \delta_n| \left| D\Phi_{s,p}(u_n)[u_n] \right| = 0,$$

thanks to the fact that by homogeneity we have

$$\left| D\Phi_{s,p}(u_n)[u_n] \right| = p |\Phi_{s,p}(u_n)|,$$

which is uniformly bounded by hypothesis (4.1). On the other hand, by weak convergence of u_n to u we also have

$$\lim_{n \rightarrow \infty} \left| D\Phi_{s,p}(u)[u_n - u] \right| = 0.$$

In conclusion we get

$$(4.3) \quad \lim_{n \rightarrow \infty} \left| D\Phi_{s,p}(u_n)[u_n - u] - D\Phi_{s,p}(u)[u_n - u] \right| = 0.$$

If we set

$$U_n(x, y) = u_n(x) - u_n(y) \quad \text{and} \quad U(x, y) = u(x) - u(y),$$

then (4.3) can be rewritten as

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left(|U_n(x, y)|^{p-2} U_n(x, y) - |U(x, y)|^{p-2} U(x, y) \right) \left(U_n(x, y) - U(x, y) \right)}{|x - y|^{N+sp}} \, dx \, dy = 0.$$

By appealing to (B.6), the previous implies for $p \geq 2$

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U_n(x, y) - U(x, y)|^p}{|x - y|^{N+sp}} \, dx \, dy = 0,$$

so that u_n strongly converges in $\widetilde{W}_0^{s,p}(\Omega)$ to u (up to a subsequence). For the case $1 < p < 2$, we use (B.5) raised to the power $p/2$, so that we get

$$\begin{aligned} & \lim_{n \rightarrow \infty} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U_n(x, y) - U(x, y)|^p}{|x - y|^{N+sp}} \, dx \, dy \\ & \leq \left(\frac{1}{p-1} \right)^{\frac{p}{2}} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{\left(|U_n(x, y)|^{p-2} U_n(x, y) - |U(x, y)|^{p-2} U(x, y) \right)^{\frac{p}{2}}}{|x - y|^{N+sp}} \\ & \quad \times \left(U_n(x, y) - U(x, y) \right)^{\frac{p}{2}} \left(|U_n(x, y)|^2 + |U(x, y)|^2 \right)^{\frac{2-p}{2} \frac{p}{2}} \, dx \, dy. \end{aligned}$$

It is now sufficient to use Hölder inequality with exponents $2/p$ and $2/(2-p)$ in order to infer again the strong convergence.

Part 2: $\lambda_2(\Omega) > \lambda_1(\Omega)$. We can use a simple topological argument, like in [2, Theorem 4.2]. Let us argue by contradiction and suppose that

$$\lambda_2(\Omega) = \inf_{f \in \mathcal{C}_1(\Omega)} \max_{u \in \text{Im}(f)} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)} = \lambda_1(\Omega),$$

so that, for all $n \in \mathbb{N}$ there exists an odd continuous mapping $f_n : \mathbb{S}^1 \rightarrow \mathcal{S}_p(\Omega)$ such that

$$(4.4) \quad \max_{u \in f_n(\mathbb{S}^1)} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)} \leq \lambda_1(\Omega) + \frac{1}{n}.$$

Let us denote by u_1 the unique (modulo the choice of the sign) global minimizer belonging to $\mathcal{S}_p(\Omega)$. Let $0 < \varepsilon \ll 1$ and consider the two neighborhoods

$$\mathcal{B}_\varepsilon^+ = \{u \in \mathcal{S}_p(\Omega) : \|u - u_1\|_{L^p(\Omega)} < \varepsilon\} \quad \mathcal{B}_\varepsilon^- = \{u \in \mathcal{S}_p(\Omega) : \|u - (-u_1)\|_{L^p(\Omega)} < \varepsilon\},$$

which are disjoint, by construction. Since the mapping f_n is odd and continuous, for every $n \in \mathbb{N}$ the image $f_n(\mathbb{S}^1)$ is symmetric and connected, then it can not be contained in $\mathcal{B}_\varepsilon^+ \cup \mathcal{B}_\varepsilon^-$, the latter being symmetric and disconnected. So we can pick an element

$$(4.5) \quad u_n \in f_n(\mathbb{S}^1) \setminus (\mathcal{B}_\varepsilon^+ \cup \mathcal{B}_\varepsilon^-).$$

This yields a sequence $\{u_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_p(\Omega)$, which is bounded in $\widetilde{W}_0^{s,p}(\Omega)$ by (4.4). Hence, there exists a function $v \in \mathcal{S}_p(\Omega)$ such that $\{u_n\}_{n \in \mathbb{N}}$ converges to v weakly in $\widetilde{W}_0^{s,p}(\Omega)$ and strongly in $L^p(\Omega)$, possibly by passing to a subsequence. By the weak convergence it follows that

$$\|v\|_{\widetilde{W}_0^{s,p}(\Omega)} \leq \liminf_{n \rightarrow \infty} \|u_n\|_{\widetilde{W}_0^{s,p}(\Omega)} = \lambda_1(\Omega).$$

This in turn shows that $v \in \mathcal{S}_p(\Omega)$ is a global minimizer, so that either $v = u_1$ or $v = -u_1$. On the other hand, by strong L^p convergence we also have

$$v \in \mathcal{S}_p(\Omega) \setminus (\mathcal{B}_\varepsilon^+ \cup \mathcal{B}_\varepsilon^-).$$

This gives a contradiction and thus $\lambda_2(\Omega) > \lambda_1(\Omega)$.

Part 3: $\lambda_2(\Omega)$ admits only sign-changing eigenfunctions. This follows directly from the previous step and Theorem 2.8. \square

The following result justifies the notation we used for $\lambda_2(\Omega)$. Indeed, *the latter is exactly the second (s, p) -eigenvalue.*

Proposition 4.2. *Let $0 < s < 1$ and $1 < p < \infty$. Let $\Omega \subset \mathbb{R}^N$ be an open and bounded set. For every eigenvalue $\lambda > \lambda_1(\Omega)$ we have*

$$\lambda_2(\Omega) \leq \lambda.$$

In particular $\lambda_1(\Omega)$ is isolated.

Proof. Let (u, λ) be an eigenpair, with $\lambda > \lambda_1(\Omega)$. Then u has to change sign in Ω , i.e. $u = u_+ - u_-$ with $u_+ \not\equiv 0$ and $u_- \not\equiv 0$, both being positive. By testing the equation solved by u against u_+ and u_- , we get

$$\lambda \int_{\Omega} u_+^p dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (u_+(x) - u_+(y)) dx dy,$$

and

$$-\lambda \int_{\Omega} u_-^p dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (u_-(x) - u_-(y)) dx dy.$$

Let us set for simplicity

$$U(x, y) = u_+(x) - u_+(y) \quad \text{and} \quad V(x, y) = u_-(x) - u_-(y),$$

then

$$u(x) - u(y) = (u_+(x) - u_-(x)) - (u_+(y) - u_-(y)) = U(x, y) - V(x, y).$$

In this way we can rewrite

$$\lambda \int_{\Omega} u_+^p dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U - V|^{p-2} (U - V)}{|x - y|^{N+sp}} U dx dy,$$

and

$$-\lambda \int_{\Omega} u_-^p dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U - V|^{p-2} (U - V)}{|x - y|^{N+sp}} V dx dy.$$

Let $(\omega_1, \omega_2) \in \mathbb{S}^1$, by multiplying the previous two identities by $|\omega_1|^p$ and $|\omega_2|^p$ and subtracting them, we can then arrive at

$$(4.6) \quad \lambda = \frac{\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{[|\omega_1|^p |U - V|^{p-2} (U - V) U - |\omega_2|^p |U - V|^{p-2} (U - V) V]}{|x - y|^{N+sp}} dx dy}{|\omega_1|^p \int_{\Omega} u_+^p dx + |\omega_2|^p \int_{\Omega} u_-^p dx}.$$

Observe that the integrand in the numerator can be written as

$$|\omega_1 U - \omega_1 V|^{p-2} (\omega_1 U - \omega_1 V) \omega_1 U - |\omega_2 U - \omega_2 V|^{p-2} (\omega_2 U - \omega_2 V) \omega_2 V.$$

We now claim that

$$(4.7) \quad \begin{aligned} & |\omega_1 U - \omega_1 V|^{p-2} (\omega_1 U - \omega_1 V) \omega_1 U - |\omega_2 U - \omega_2 V|^{p-2} (\omega_2 U - \omega_2 V) \omega_2 V \\ & \geq |\omega_1 U - \omega_2 V|^p. \end{aligned}$$

Let us now assume (4.7) for the moment and show how the proof ends. Indeed, if we define the following element of $\mathcal{C}_1(\Omega)$

$$f(\omega) = \frac{\omega_1 u_+ - \omega_2 u_-}{\left(|\omega_1|^p \int_{\Omega} u_+^p dx + |\omega_2|^p \int_{\Omega} u_-^p dx \right)^{1/p}}, \quad \omega = (\omega_1, \omega_2) \in \mathbb{S}^1,$$

then with the notations above we have

$$\|f(\omega)\|_{\widetilde{W}_0^{s,p}(\Omega)}^p = \frac{\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\omega_1 U - \omega_2 V|^p}{|x - y|^{N+sp}} dx dy}{|\omega_1|^p \int_{\Omega} u_+^p dx + |\omega_2|^p \int_{\Omega} u_-^p dx}.$$

If we now use the pointwise inequality (4.7) and recall the relation (4.6) for λ , we get

$$\|f(\omega)\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq \lambda, \quad \text{for every } \omega \in \mathbb{S}^1.$$

By appealing to the definition of $\lambda_2(\Omega)$ we get the desired conclusion. Observe that $\lambda_1(\Omega)$ is isolated thanks to the fact that $\lambda_2(\Omega) > \lambda_1(\Omega)$.

In order to conclude the proof, let us now prove the vital pointwise inequality (4.7). We start observing that $U \cdot V \leq 0$.

If $\omega_1 = 0$, then the left-hand side of (4.7) reduces to

$$-|\omega_2 U - \omega_2 V|^{p-2} (\omega_2 U - \omega_2 V) \omega_2 V = -|\omega_2|^p |U - V|^{p-2} (U - V) V.$$

If $V = 0$, the inequality is obvious. If $V < 0$, then $U \geq 0$ so that $U - V \geq -V$. By using the monotonicity of the map $\tau \mapsto |\tau|^{p-2} \tau$ we get

$$-|\omega_2|^p |U - V|^{p-2} (U - V) V \geq -|\omega_2|^p |-V|^{p-2} (-V) V = |\omega_2 V|^p,$$

which coincides with the right-hand side of (4.7) when $\omega_1 = 0$. If on the contrary we have $V > 0$, then $U \leq 0$ and in this case $U - V \leq -V$. Using again the monotonicity of the map $\tau \mapsto |\tau|^{p-2} \tau$, we get again

$$-|\omega_2|^p |U - V|^{p-2} (U - V) V \geq -|\omega_2|^p |-V|^{p-2} (-V) V = |\omega_2|^p V^p.$$

This proves (4.7) for the case $\omega_1 = 0$.

We now assume $\omega_1 \neq 0$, then by dividing everything by $|\omega_1|^p$ we get that (4.7) is equivalent to prove

$$|U - V|^{p-2} (U - V) C \geq \left| \frac{\omega_2}{\omega_1} U - \frac{\omega_2}{\omega_1} V \right|^{p-2} \left(\frac{\omega_2}{\omega_1} U - \frac{\omega_2}{\omega_1} V \right) \frac{\omega_2}{\omega_1} V + \left| U - \frac{\omega_2}{\omega_1} V \right|^p.$$

We now observe that the previous inequality is a direct consequence of Lemma B.1, thus the proof is complete. \square

Remark 4.3. The fact that $\lambda_1(\Omega)$ is isolated was also proved in [21, Theorem 19], under the restriction $sp > N$. The restriction was needed in order to have the eigenfunctions continuous, a fact that we would have now for free from Corollary 3.14. However, our proof is different and does not need continuity of eigenfunctions.

Remark 4.4. It is not difficult to see that $\lambda_2(\Omega)$ coincides with λ_2 defined in [18, Section 2] by means of a cohomological index. We leave the details to the interested reader.

5. MOUNTAIN PASS CHARACTERIZATION

In this section we prove an alternative characterization of $\lambda_2(\Omega)$ as a mountain pass level. In order to prove such a characterization, the following technical result will be useful.

Lemma 5.1. *Let $1 < p < \infty$ and $0 < s < 1$. Let $\Omega \subset \mathbb{R}^N$ be an open and bounded set. For every $u \in \mathcal{S}_p(\Omega)$ we set*

$$U(x, y) = u_+(x) - u_+(y) \quad \text{and} \quad V(x, y) = u_-(x) - u_-(y),$$

and we define the continuous curve on $\mathcal{S}_p(\Omega)$

$$\gamma_t = \frac{u_+ - \cos(\pi t) u_-}{\|u_+ - \cos(\pi t) u_-\|_{L^p(\Omega)}}, \quad t \in \left[0, \frac{1}{2}\right].$$

Let us suppose that we have

$$(5.1) \quad \begin{aligned} & \left(\int_{\Omega} u_-^p dx \right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U - V|^{p-2} (U - V) U}{|x - y|^{N+sp}} dx dy \\ & + \left(\int_{\Omega} u_+^p dx \right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U - V|^{p-2} (U - V) V}{|x - y|^{N+sp}} dx dy \leq 0, \end{aligned}$$

then there holds

$$\|\gamma_t\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p, \quad t \in \left[0, \frac{1}{2}\right].$$

Proof. We start by observing that

$$\|\gamma_t\|_{\widetilde{W}_0^{s,p}(\Omega)}^p = \frac{\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U - \cos(\pi t) V|^p}{|x - y|^{N+sp}} dx dy}{\int_{\Omega} u_+^p dx + |\cos(\pi t)|^p \int_{\Omega} u_-^p dx}, \quad t \in \left[0, \frac{1}{2}\right].$$

By definition we have $U \cdot V \leq 0$, then by using Lemma B.1 we get

$$\begin{aligned} \|\gamma_t\|_{\widetilde{W}_0^{s,p}(\Omega)}^p &\leq \frac{\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) U}{|x-y|^{N+sp}} dx dy}{\int_{\Omega} u_+^p dx + |\cos(\pi t)|^p \int_{\Omega} u_-^p dx} \\ &\quad - |\cos(\pi t)|^p \frac{\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) V}{|x-y|^{N+sp}} dx dy}{\int_{\Omega} u_+^p dx + |\cos(\pi t)|^p \int_{\Omega} u_-^p dx}, \quad t \in \left[0, \frac{1}{2}\right]. \end{aligned}$$

Observe that the term on the right-hand side has the form

$$\frac{a-sb}{c+sd}, \quad \text{for } s \in [0, 1].$$

In order to get the conclusion, it is then sufficient to observe that the function

$$s \mapsto \frac{a-sb}{c+sd},$$

is monotone increasing if and only if

$$-cb - da \geq 0,$$

that is if and only if

$$\begin{aligned} &\left(\int_{\Omega} u_-^p dx\right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) U}{|x-y|^{N+sp}} dx dy \\ &\quad + \left(\int_{\Omega} u_+^p dx\right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) V}{|x-y|^{N+sp}} dx dy \leq 0, \end{aligned}$$

which is exactly hypothesis (5.1). By using this and recalling that u has unit L^p norm, we get

$$\begin{aligned} \|\gamma_t\|_{\widetilde{W}_0^{s,p}(\Omega)}^p &\leq \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) U}{|x-y|^{N+sp}} dx dy \\ &\quad - \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) V}{|x-y|^{N+sp}} dx dy = \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p, \quad t \in \left[0, \frac{1}{2}\right], \end{aligned}$$

as desired. \square

The following simple remark will be important.

Remark 5.2. Let $u \in \mathcal{S}_p(\Omega)$ be a function that does not satisfy (5.1), i.e.

$$\begin{aligned} &\left(\int_{\Omega} u_-^p dx\right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) U}{|x-y|^{N+sp}} dx dy \\ &\quad + \left(\int_{\Omega} u_+^p dx\right) \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|U-V|^{p-2} (U-V) V}{|x-y|^{N+sp}} dx dy > 0 \end{aligned}$$

Then it is easy to see that the function $v = -u$ satisfies (5.1).

Let us define

$$\Gamma(u_1, -u_1) = \{\gamma \in C^0([0, 1]; \mathcal{S}_p(\Omega)) : \gamma_0 = u_1, \gamma_1 = -u_1\},$$

the set of continuous curves on $\mathcal{S}_p(\Omega)$ connecting the two global minimizers u_1 and $-u_1$. We have the following characterization for $\lambda_2(\Omega)$. The idea of the proof is the same as in [2, Proposition 5.4].

Theorem 5.3 (Mountain pass characterization). *Let $1 < p < \infty$ and $0 < s < 1$. Let $\Omega \subset \mathbb{R}^N$ be an open and bounded set, then we have*

$$\lambda_2(\Omega) = \inf_{\gamma \in \Gamma(u_1, -u_1)} \max_{u \in \text{Im}(\gamma)} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p.$$

Proof. We first observe that the inequality

$$\lambda_2(\Omega) \leq \inf_{\gamma \in \Gamma(u_1, -u_1)} \max_{u \in \gamma} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p,$$

is easily seen to hold. Indeed, given $\gamma \in \Gamma(u_1, -u_1)$, we can consider the closed loop $\{\gamma\} \cup \{-\gamma\}$ and identify it with an element of $\mathcal{C}_1(\Omega)$.

Let us now prove the converse. For every $n \in \mathbb{N}$, we pick $f_n \in \mathcal{C}_1(\Omega)$ “almost optimal”, i.e. such that

$$\max_{u \in \text{Im}(f_n)} \|u\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq \lambda_2(\Omega) + \frac{1}{n}.$$

Since f_n is odd, the set $\text{Im}(f_n)$ is symmetric with respect to the origin. Then thanks to Remark 5.2 there exists $u_n \in \text{Im}(f_n)$ which verifies hypothesis (5.1). This implies that on the curve

$$\gamma_{n,t} = \frac{(u_n)_+ - \cos(\pi t) (u_n)_-}{\|(u_n)_+ - \cos(\pi t) (u_n)_-\|_{L^p(\Omega)}}, \quad t \in \left[0, \frac{1}{2}\right],$$

we have

$$(5.2) \quad \|\gamma_{n,t}\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq \lambda_2(\Omega) + \frac{1}{n}, \quad t \in \left[0, \frac{1}{2}\right].$$

Observe that the curve γ_n is connecting u_n to its (renormalized in L^p) positive part, without increasing the energy. We now in turn connect the function $(u_n)_+ / \|(u_n)_+\|_{L^p(\Omega)}$ to the first eigenfunction u_1 : at this aim, we recall that on the curve

$$\sigma_{n,t} = \left((1-t) \frac{(u_n)_+^p}{\|(u_n)_+\|_{L^p(\Omega)}^p} + t u_1^p \right)^{\frac{1}{p}}, \quad t \in [0, 1],$$

our energy functional is convex (see [14, Lemma 4.1] or also [1, Proposition 4.1]), i.e.

$$\|\sigma_{n,t}\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq (1-t) \frac{\|(u_n)_+\|_{\widetilde{W}_0^{s,p}(\Omega)}^p}{\|(u_n)_+\|_{L^p(\Omega)}^p} + t \|u_1\|_{\widetilde{W}_0^{s,p}(\Omega)}^p.$$

Thus in particular from (5.2) we have

$$\|\sigma_{n,t}\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq \lambda_2(\Omega) + \frac{1}{n}, \quad t \in [0, 1].$$

Thus we can glue together γ_n and σ_n and build the new curve

$$\widetilde{\gamma}_{n,t} = \begin{cases} \gamma_{n,t}, & t \in [0, 1/2], \\ \sigma_{n,(2t-1)}, & t \in [1/2, 1] \end{cases}$$

which is connecting u_n to u_1 and on which the energy is always less than $\lambda_2(\Omega) + 1/n$. Finally, if we glue together $\widetilde{\gamma}_n$, $-\widetilde{\gamma}_n$ and f_n , perform a suitable reparameterization and use the fact that the energy functional is even, we obtain a new continuous curve $\Sigma_n \in \Gamma(u_1, -u_1)$ such that

$$\max_{t \in [0,1]} \|\Sigma_{n,t}\|_{\widetilde{W}_0^{s,p}(\Omega)}^p \leq \lambda_2(\Omega) + \frac{1}{n}, \quad n \in \mathbb{N}.$$

This of course implies that

$$\inf_{\gamma \in \Gamma(u_1, -u_1)} \max_{u \in \text{Im}(\gamma)} \|u\|_{\widehat{W}_0^{s,p}(\Omega)}^p \leq \lambda_2(\Omega) + \frac{1}{n}.$$

By taking the limit as n goes to ∞ , we finally obtain the desired conclusion. \square

Remark 5.4. In the nonlocal case, the previous characterization has been proven for $p = 2$ in [16], in the case of the Fučík spectrum, by adapting the proof of [7].

6. A SHARP LOWER BOUND

We are going to prove a sharp lower bound on $\lambda_2(\Omega)$ in terms of the measure of Ω . The following simple result will be important.

Lemma 6.1 (Nodal domains). *Let $\lambda > \lambda_1(\Omega)$ be an (s, p) -eigenvalue. Let $u \in \mathcal{S}_p(\Omega)$ be an associated eigenfunction and set*

$$\Omega_+ = \{x \in \Omega : u(x) > 0\} \quad \text{and} \quad \Omega_- = \{x \in \Omega : u(x) < 0\}.$$

Then we have

$$\lambda > \max\{\lambda_1(\Omega_+), \lambda_1(\Omega_-)\}.$$

Proof. We first observe that since (s, p) -eigenfunctions are continuous by Corollary 3.14, the sets Ω_+ and Ω_- are open. Thus $\lambda_1(\Omega_+)$ and $\lambda_1(\Omega_-)$ are well-defined.

The case $p \geq 2$ is already contained in [21, Theorem 17], so let us focus on the case $1 < p < 2$. Since u is sign-changing, we can write $u = u_+ - u_-$, where u_+ and u_- are the positive and negative parts respectively. By testing the equation solved by u against u_+ we get

$$\lambda \int_{\Omega} |u_+|^p dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(x) - u(y)|^{p-2} (u(x) - u(y))}{|x - y|^{N+sp}} (u_+(x) - u_+(y)) dx dy.$$

Then we can apply Lemma B.2 with the choices

$$a = u_+(x) - u_+(y) \quad \text{and} \quad b = u_-(x) - u_-(y),$$

and obtain

$$\lambda \int_{\Omega} |u_+|^p dx > \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u_+(x) - u_+(y)|^p}{|x - y|^{N+sp}} dx dy.$$

Since u_+ is admissible for the variational problem defining $\lambda_1(\Omega_+)$, we get $\lambda > \lambda_1(\Omega_+)$.

For the other set Ω_- , we proceed similarly by testing the equation against u_- , thus getting

$$\lambda \int_{\Omega} |u_-|^p dx = \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|u(y) - u(x)|^{p-2} (u(y) - u(x))}{|x - y|^{N+sp}} (u_-(x) - u_-(y)) dx dy.$$

If we now use again Lemma B.2, this time with the choices

$$a = u_-(x) - u_-(y) \quad \text{and} \quad b = u_+(x) - u_+(y),$$

we get also the estimate $\lambda > \lambda_1(\Omega_-)$. This concludes the proof. \square

The following is the main result of this section. This is the nonlocal version of the so-called *Hong-Krahn-Szego inequality* (see [3, Theorem 3.2]), which in the local case asserts that the second eigenvalue of the Dirichlet-Laplacian is minimized by the disjoint union of two equal balls, among sets of given measure.

Theorem 6.2 (Nonlocal Hong-Krahn-Szego inequality). *Let $0 < s < 1$ and $1 < p < \infty$. For every $\Omega \subset \mathbb{R}^N$ open and bounded set we have*

$$(6.1) \quad \lambda_2(\Omega) > \lambda_1(B),$$

where B is any N -dimensional ball such that $|B| = |\Omega|/2$. Equality is never attained in (6.1), but the estimate is sharp in the following sense: if $\{x_n\}_{n \in \mathbb{N}}, \{y_n\}_{n \in \mathbb{N}} \subset \mathbb{R}^N$ are such that

$$\lim_{n \rightarrow \infty} |x_n - y_n| = +\infty,$$

and we define $\Omega_n := B_R(x_n) \cup B_R(y_n)$, then

$$\lim_{n \rightarrow \infty} \lambda_2(\Omega_n) = \lambda_1(B_R).$$

Proof. We divide the proof in two steps: at first we prove (6.1), then we prove its sharpness.

Inequality. Let $u \in \mathcal{S}_p(\Omega)$ be an eigenfunction associated to $\lambda_2(\Omega)$. By Theorem 4.1, we know that u is sign-changing, thus we define

$$\Omega_+ := \{x \in \Omega : u(x) > 0\} \quad \text{and} \quad \Omega_- := \{x \in \Omega : u(x) < 0\}.$$

By Lemma 6.1 and the *nonlocal Faber-Krahn inequality* (see [4, Theorem 3.5]), we have

$$\lambda_2(\Omega) > \lambda_1(\Omega_+) \geq \lambda_1(B_{R_1}) \quad \text{and} \quad \lambda_2(\Omega) > \lambda_1(\Omega_-) \geq \lambda_1(B_{R_2}).$$

where B_{R_1} and B_{R_2} are such that $|B_{R_1}| = |\Omega_+|$ and $|B_{R_2}| = |\Omega_-|$. Thus

$$(6.2) \quad \lambda_2(\Omega) > \max\{\lambda_1(B_{R_1}), \lambda_1(B_{R_2})\}.$$

By the scaling properties of λ_1 we have $\lambda_1(B_R) = R^{-s/p} \lambda_1(B_1)$, moreover we have the constraint

$$|B_{R_1}| + |B_{R_2}| \leq |\Omega|.$$

Then it is easy to see that the right-hand side of (6.2) is minimized when $|B_{R_1}| = |B_{R_2}| = |\Omega|/2$, which implies the desired estimate (6.1).

Sharpness. In order to prove the second part of the claim, we define

$$\Omega_n := B_R(x_n) \cup B_R(y_n),$$

where $\{x_n\}, \{y_n\} \subset \mathbb{R}^N$ are such that $|x_n - y_n|$ diverges as n goes to ∞ . Thus we can suppose that the two balls are disjoint. Let u and v be the positive normalized first eigenfunctions on $B_R(x_n)$ and $B_R(y_n)$ respectively (observe that their shape does not depend on the center of the ball), then we set for simplicity

$$a(x, y) = u(x) - u(y) \quad \text{and} \quad b(x, y) = v(x) - v(y).$$

By Lemma B.3 and the definition of $\lambda_2(\Omega)$ we have⁸

$$\begin{aligned} \lambda_2(\Omega_n) &\leq \max_{|\omega_1|^p + |\omega_2|^p = 1} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\omega_1 a - \omega_2 b|^p}{|x - y|^{N+sp}} dx dy \\ &\leq \max_{|\omega_1|^p + |\omega_2|^p = 1} \left[\int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\omega_1|^p |a|^p}{|x - y|^{N+sp}} dx dy + \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{|\omega_2|^p |b|^p}{|x - y|^{N+sp}} dx dy \right. \\ &\quad \left. + c_p \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(|\omega_1 a|^2 + |\omega_2 b|^2)^{\frac{p-2}{2}} |\omega_1 \omega_2 a b|}{|x - y|^{N+sp}} dx dy \right] \\ &= \lambda_1(B_R) + c_p \max_{|\omega_1|^p + |\omega_2|^p = 1} \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{(|\omega_1 a|^2 + |\omega_2 b|^2)^{\frac{p-2}{2}} |\omega_1 \omega_2 a b|}{|x - y|^{N+sp}} dx dy. \end{aligned}$$

⁸For simplicity, we used the change of variable $(\omega_1, \omega_2) \mapsto (|\omega_1|^{\frac{2-p}{p}} \omega_1, |\omega_2|^{\frac{2-p}{p}} \omega_2)$.

Observe that, since

$$a b = -u(x) v(y) - u(y) v(x),$$

the numerator in the last integral is nonzero only if $(x, y) \in B_R(x_n) \times B_R(y_n)$ or $(x, y) \in B_R(y_n) \times B_R(x_n)$. We set

$$\mathcal{R} := 2 \max_{|\omega_1|^p + |\omega_2|^p = 1} \int_{B_R(x_n)} \int_{B_R(y_n)} (|\omega_1 a|^2 + |\omega_2 b|^2)^{\frac{p-2}{2}} |\omega_1 \omega_2 a b| dx dy < \infty,$$

therefore we have

$$\lim_{n \rightarrow \infty} \lambda_2(\Omega_n) \leq \lambda_1(B_R) + \lim_{n \rightarrow \infty} \frac{c_p \mathcal{R}}{(|x_n - y_n| - 2R)^{N+sp}} = \lambda_1(B_R).$$

This concludes the proof. \square

Remark 6.3. The previous result can be stated in scaling invariant form as follows

$$(6.3) \quad \lambda_2(\Omega) > \left(\frac{2|B|}{|\Omega|} \right)^{\frac{sp}{N}} \lambda_1(B),$$

where B is any N -dimensional ball.

APPENDIX A. SOME USEFUL INEQUALITIES I

We will repeatedly use that for $1 < p < \infty$ the real function

$$J_p(t) := |t|^{p-2} t,$$

is monotone increasing.

Lemma A.1 (Towards subsolutions). *Let $1 < p < \infty$ and $f : \mathbb{R} \rightarrow \mathbb{R}$ be a convex function, then*

$$(A.1) \quad \begin{aligned} & |a - b|^{p-2} (a - b) \left[A |f'(a)|^{p-2} f'(a) - B |f'(b)|^{p-2} f'(b) \right] \\ & \geq |f(a) - f(b)|^{p-2} (f(a) - f(b)) (A - B), \end{aligned}$$

for every $a, b \in \mathbb{R}$ and every $A, B \geq 0$.

Proof. By convexity of f we have

$$(A.2) \quad f(b) - f(a) \geq f'(a) (b - a) \quad \text{and} \quad f(a) - f(b) \geq f'(b) (a - b).$$

By writing the left-hand side of (A.1) as

$$\begin{aligned} & |a - b|^{p-2} (a - b) \left[A |f'(a)|^{p-2} f'(a) - B |f'(b)|^{p-2} f'(b) \right] \\ & = |a - b|^{p-2} (a - b) A |f'(a)|^{p-2} f'(a) - |a - b|^{p-2} (a - b) B |f'(b)|^{p-2} f'(b) \\ & = J_p(f'(a) (a - b)) A - J_p(f'(b) (a - b)) B, \end{aligned}$$

we can get the conclusion by simply using (A.2) and the monotonicity of J_p . \square

The next pointwise inequality generalizes a similar estimate in [4, Appendix C].

Lemma A.2 (Towards Moser's iteration). *Let $1 < p < \infty$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function. We define*

$$G(t) = \int_0^t g'(\tau)^{\frac{1}{p}} d\tau, \quad s \in \mathbb{R},$$

then we have

$$(A.3) \quad |a - b|^{p-2} (a - b) (g(a) - g(b)) \geq |G(a) - G(b)|^p.$$

Proof. We first observe that we can suppose $a > b$ without loss of generality. Then

$$\begin{aligned} |a - b|^{p-2} (a - b) (g(a) - g(b)) &= (a - b)^{p-1} (g(a) - g(b)) = (a - b)^{p-1} \int_b^a g'(\tau) d\tau \\ &= (a - b)^{p-1} \int_b^a G'(\tau)^p d\tau \geq \left(\int_b^a G'(\tau) d\tau \right)^p, \end{aligned}$$

thanks to Jensen inequality. \square

Remark A.3. With the same proof one can show that if g is decreasing, then

$$|a - b|^{p-2} (a - b) (g(b) - g(a)) \geq |H(a) - H(b)|^p, \quad \text{where} \quad H(t) = \int_0^t -g'(\tau)^{\frac{1}{p}} d\tau.$$

Lemma A.4. Let $\beta \geq 1$, then for every $a, b \geq 0$ we have

$$(A.4) \quad |a - b|^p \left(a^{\beta-1} + b^{\beta-1} \right) \leq \left(\max\{1, (3 - \beta)\} \right) |a - b|^{p-2} (a - b) (a^\beta - b^\beta).$$

Proof. We first observe that (A.4) is trivially true for $a = b$, thus let us consider $a \neq b$. It is not restrictive to assume that $a > b$, then (A.4) is equivalent to

$$(1 - t)^p \left(1 + t^{\beta-1} \right) \leq C (1 - t)^{p-1} (1 - t^\beta), \quad \text{for } 0 \leq t < 1,$$

that is

$$(A.5) \quad (1 - t) \left(1 + t^{\beta-1} \right) \leq C (1 - t^\beta).$$

By observing that

$$(1 - t) \left(1 + t^{\beta-1} \right) = (1 - t^\beta) + t^{\beta-1} - t,$$

and remembering that $0 \leq t < 1$, we easily get the conclusion for $\beta \geq 2$, since $t^{\beta-1} - t \leq 0$ in this case. If on the contrary $1 < \beta < 2$, then by concavity of the function $\tau \mapsto \tau^{\beta-1}$ we have

$$\begin{aligned} t^{\beta-1} - t &= (t^{\beta-1} - 1) - (t - 1) \leq (\beta - 1) (t - 1) - (t - 1) = (2 - \beta) (1 - t) \\ &\leq (2 - \beta) (1 - t^\beta). \end{aligned}$$

This finally shows (A.4) for $1 < \beta < 2$ as well. \square

Lemma A.5. Let $p \geq 1$, then

$$\left(\frac{1}{\beta} \right)^{\frac{1}{p}} \frac{\beta + p - 1}{p} \geq 1, \quad \text{for every } \beta > 0.$$

Proof. For $p = 1$ there is nothing to prove, thus let us assume that $p > 1$. The result follows from convexity of the function $t \mapsto t^p$, which implies

$$\beta - 1 \geq p \left(\beta^{\frac{1}{p}} - 1 \right).$$

By adding p on both sides, we get the conclusion. \square

APPENDIX B. SOME USEFUL INEQUALITIES II

We still use the notation $J_p(t) = |t|^{p-2} t$.

Lemma B.1. *Let $1 < p < \infty$ and $U, V \in \mathbb{R}$ such that $UV \leq 0$. We define the following function*

$$g(t) = |U - tV|^p + |U - V|^{p-2} (U - V) V |t|^p, \quad t \in \mathbb{R}.$$

Then we have

$$g(t) \leq g(1) = |U - V|^{p-2} (U - V) U, \quad t \in \mathbb{R}.$$

Proof. Let us start observing that if $U = 0$, then we have

$$g(t) = 0, \quad \text{for every } t \in \mathbb{R}.$$

In the same manner, if $V = 0$, then we have

$$g(t) = |U|^p, \quad \text{for every } t \in \mathbb{R}.$$

In both cases, the conclusion trivially holds true.

Thus we can suppose that $UV \neq 0$. Then we have

$$\begin{aligned} g'(t) &= -p|U - tV|^{p-2} (U - tV) V + p|U - V|^{p-2} (U - V) V |t|^{p-2} t \\ &= pV \left[|t(U - V)|^{p-2} t(U - V) - |U - tV|^{p-2} (U - tV) \right] \\ &= pV \left[J_p(tU - tV) - J_p(U - tV) \right]. \end{aligned}$$

We distinguish two cases.

(1) *Case $V < 0$ and $U > 0$:* then we have

$$g'(t) \geq 0 \quad \Longleftrightarrow \quad t(U - V) \leq U - tV \quad \Longleftrightarrow \quad t \leq 1.$$

This implies that $t = 1$ is a global maximum point for the function g .

(2) *Case $V > 0$ and $U < 0$:* we now have

$$g'(t) \geq 0 \quad \Longleftrightarrow \quad t(U - V) \geq U - tV \quad \Longleftrightarrow \quad t \leq 1,$$

since now $C < 0$. Again, we get that $t = 1$ is a global maximum point.

In both cases, we get the desired conclusion. \square

Lemma B.2. *Let $1 < p < \infty$, then for every $a, b \in \mathbb{R}$ such that $ab \leq 0$, we have*

$$(B.1) \quad |a - b|^{p-2} (a - b) a \geq \begin{cases} |a|^p - (p-1) |a - b|^{p-2} ab, & \text{if } 1 < p \leq 2, \\ |a|^p - (p-1) |a|^{p-2} ab, & \text{if } p > 2. \end{cases}.$$

Proof. We start with some elementary considerations. First of all, there is no loss of generality in supposing $a \geq 0$ and $b \leq 0$. Then we notice that the function J_p on $[0, \infty)$ is convex for $p > 2$ and concave for $1 < p \leq 2$. Thus

$$(B.2) \quad J_p(x) + J'_p(y) (y - x) \leq J_p(y), \quad \text{for } 1 < p \leq 2, \quad 0 \leq x \leq y,$$

and

$$(B.3) \quad J_p(x) + J'_p(x) (y - x) \leq J_p(y), \quad \text{for } p \geq 2, \quad 0 \leq x \leq y.$$

We now come to the proof of (B.1), starting with the case $1 < p \leq 2$. By using (B.2) with the choices

$$y = a - b \quad \text{and} \quad x = a,$$

we get

$$\begin{aligned} |a - b|^{p-2} (a - b) &= J_p(a - b) \geq J_p(a) - J'_p(a - b) b \\ &= |a|^{p-2} a - (p - 1) |a - b|^{p-2} b. \end{aligned}$$

and multiplying by $a \geq 0$ we conclude. This ends the proof in the case $1 < p \leq 2$.

The case $p > 2$ is handled in a similar manner, by using (B.3) instead of (B.2). \square

Lemma B.3. *Let $1 < p < \infty$. Then there exists $c_p > 0$ such that for every $a, b \in \mathbb{R}$ we have*

$$(B.4) \quad |a - b|^p \leq |a|^p + |b|^p + c_p (|a|^2 + |b|^2)^{\frac{p-2}{2}} |a b|.$$

Proof. We first suppose $a b \geq 0$, without loss of generality we can suppose that $a, b \geq 0$ and $a \geq b$. Then we have

$$|a - b|^p = (a - b)^p \leq a^p \leq |a|^p + |b|^p,$$

thus (B.4) is proved.

Let us consider now the case $a b \leq 0$. Without loss of generality, we can suppose that $a \geq 0$ and $b \leq 0$. Then (B.4) is equivalent to

$$(a + \tau)^p \leq a^p + \tau^p + c_p (a^2 + \tau^2)^{\frac{p-2}{2}} a \cdot \tau, \quad a, \tau \geq 0.$$

Of course this is easily seen to be true if $a = 0$, so let us take $a > 0$ and divide the previous by a^p . Then we are reduced to show that

$$(1 + m)^p \leq 1 + m^p + c_p (1 + m^2)^{\frac{p-2}{2}} m,$$

that is

$$\sup_{m>0} \frac{(1 + m)^p - 1 - m^p}{m (1 + m^2)^{\frac{p-2}{2}}} < +\infty.$$

To this end, it is sufficient to observe that

$$\lim_{m \rightarrow 0^+} \frac{(1 + m)^p - 1 - m^p}{m (1 + m^2)^{\frac{p-2}{2}}} = p \quad \text{and} \quad \lim_{m \rightarrow +\infty} \frac{(1 + m)^p - 1 - m^p}{m (1 + m^2)^{\frac{p-2}{2}}} = p.$$

This concludes the proof. \square

Finally, we recall the following classical inequalities.

Lemma B.4. *For $1 < p \leq 2$, we have*

$$(B.5) \quad (|b|^{p-2} b - |a|^{p-2} a) (b - a) \geq (p - 1) \frac{|b - a|^2}{(|b|^2 + |a|^2)^{\frac{2-p}{2}}}, \quad a, b \in \mathbb{R}.$$

For $2 < p < \infty$ we have

$$(B.6) \quad (|b|^{p-2} b - |a|^{p-2} a) (b - a) \geq 2^{2-p} |b - a|^p, \quad t, s \in \mathbb{R}.$$

Proof. For the first we simply observe that

$$\begin{aligned} \left(J_p(b) - J_p(a) \right) (b - a) &= (b - a) \int_0^1 \frac{d}{dt} J_p((1 - t) a + t b) dt \\ &= (p - 1) (b - a)^2 \int_0^1 |(1 - t) a + t b|^{p-2} dt, \end{aligned}$$

and then use that

$$|(1 - t) a + t b|^2 \leq (1 - t) |a|^2 + t |b|^2 \leq |a|^2 + |b|^2.$$

For the second the reader is referred to [22, Section 10]. \square

REFERENCES

- [1] L. Brasco, G. Franzina, *Convexity properties of Dirichlet integrals and Picone-type inequalities*, Kodai Math. J., **37** (2014), 769–799.
- [2] L. Brasco, G. Franzina, *An anisotropic eigenvalue problem of Stekloff type and weighted Wulff inequalities*, NoDEA Nonlinear Differential Equations and Applications, **20** (2013), 1795–1830.
- [3] L. Brasco, G. Franzina, *On the Hong-Krahn-Szego inequality for the p -Laplace operator*, Manuscripta Math., **141** (2013), 537–557.
- [4] L. Brasco, E. Lindgren, E. Parini, *The fractional Cheeger problem*, Interfaces Free Bound., **16** (2014), 419–458.
- [5] L. Caffarelli, L. Silvestre, *An extension problem related to the fractional Laplacian*, Comm. Partial Differential Equations, **32** (2007), 1245–1260.
- [6] M. Cuesta, *Minimax theorems on C^1 manifolds via Ekeland variational principle*, Abstr. Appl. Anal., **13** (2003), 757–768.
- [7] M. Cuesta, D. G. De Figueiredo, J.-P. Gossez, *The beginning of the Fučík spectrum for the p -Laplacian*, Journal of Differential Equations, **159** (1999), 212–238.
- [8] A. Di Castro, T. Kuusi, G. Palatucci, *Local behavior of fractional p -minimizers*, preprint (2014), available at <http://cvgmt.sns.it/paper/2379/>
- [9] A. Di Castro, T. Kuusi, G. Palatucci, *Nonlocal Harnack inequalities*, J. Funct. Anal., **267** (2014), 1807–1836.
- [10] E. Di Nezza, G. Palatucci, E. Valdinoci, *Hitchhikers guide to the fractional Sobolev spaces*, Bull. Sci. Math., **136**, 521–573.
- [11] B. Dyda, *A fractional order Hardy inequality*, Illinois J. Math., **48** (2004), 575–588.
- [12] P. Drábek, S. B. Robinson, *Resonance problems for the p -Laplacian*, J. Funct. Anal., **169** (1999), 189–200.
- [13] R. L. Frank, R. Seiringer, *Non-linear ground state representations and sharp Hardy inequalities*, J. Funct. Anal., **255** (2008), 3407–3430.
- [14] G. Franzina, G. Palatucci, *Fractional p -eigenvalues*, Riv. Mat. Univ. Parma, **5** (2014),
- [15] G. Franzina, E. Valdinoci, *Geometric analysis of fractional phase transition interfaces*, in “Geometric Properties for Parabolic and Elliptic PDE’s”, Springer INdAM Series Volume **2**, 2013, 117–130.
- [16] S. Goyal, K. Sreenadh, *On the Fučík spectrum of non-local elliptic operators*, to appear on NoDEA Nonlinear Differential Equations and Applications, **21** (2014), 567–588.
- [17] I. Hong, *On an inequality concerning the eigenvalue problem of membrane*, Kōdai Math. Sem. Rep., **6** (1954), 113–114.
- [18] A. Iannizzotto, M. Squassina, *Weyl-type laws for fractional p -eigenvalue problems*, Asymptotic Anal., **88** (2014), 233–245.
- [19] E. Krahn, *Über Minimaleigenschaften der Kugel in drei und mehr Dimensionen*, Acta Comm. Univ. Dorpat., **A9** (1926), 1–44.
- [20] T. Kuusi, G. Mingione, Y. Sire, *Nonlocal equations with measure data*, preprint (2014), available at <http://cvgmt.sns.it/paper/2466/>
- [21] E. Lindgren, P. Lindqvist, *Fractional eigenvalues*, Calc. Var. Partial Differential Equations, **49** (2014), 795–826.
- [22] P. Lindqvist, *Notes on the p -Laplace equation*, Report. University of Jyväskylä Department of Mathematics and Statistics, 102. University of Jyväskylä, Jyväskylä (2006).
- [23] G. Pólya, *On the characteristic frequencies of a symmetric membrane*, Math. Zeitschr., **63** (1955), 331–337.

AIX-MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 39 RUE FRÉDÉRIC JOLIOT CURIE, 13453 MARSEILLE, FRANCE

E-mail address: lorenzo.brasco@univ-amu.fr

E-mail address: enea.parini@univ-amu.fr